arXiv:1503.05880v2 [math.AG] 20 May 2016

THREE COMBINATORIAL FORMULAS FOR TYPE A QUIVER POLYNOMIALS AND K-POLYNOMIALS

RYAN KINSER, ALLEN KNUTSON, AND JENNA RAJCHGOT

Abstract. We provide combinatorial formulas for the multidegree and K-polynomial of an (arbitrarily oriented) type A quiver locus embedded inside of its representation space. These formulas are generalizations of three of Knutson-Miller-Shimozono's formulas from the equioriented setting:

- The *ratio formulas* express each K-polynomial as a ratio of specialized double Grothendieck polynomials, and each multidegree as a ratio of specialized double Schubert polynomials.
- The *pipe formulas* express each K-polynomial as an alternating sum over pipe dreams that fit inside of a particular shape, and each multidegree as a positive sum over reduced pipe dreams that fit inside of that same shape.
- The *component formulas* express each K-polynomial as an alternating sum of products of Grothendieck polynomials, and each multidegree as a positive sum of products of Schubert polynomials. The summands are indexed by lacing diagrams associated to the type A quiver locus. The K-polynomial component formula was first conjectured by Buch-Rimányi, and the multidegree component formula was first proved by Buch-Rimányi.

Contents

1. Introduction 1
2. Background and preliminary results 4
3. Bipartite ratio and pipe formulas 22
4. Degeneration and bipartite component formula 24
5. Generalizing the formulas to arbitrary orientation 33
References 38

1. Introduction

1.1. **Context.** Let Q be a quiver and $\mathbf{d}$ a dimension vector for Q. The variety $\mathtt{rep}_Q(\mathbf{d})$, a product of matrix spaces, parametrizes representations of Q of dimension vector $\mathbf{d}$ in some fixed basis. The base change group $GL(\mathbf{d})$ acts on $\mathtt{rep}_Q(\mathbf{d})$ so that orbits correspond to isomorphism classes (see §2.1). A quiver locus is, by definition, an orbit closure for this action.

Quiver loci have been studied in representation theory of finite-dimensional algebras since at least the early 1980s, with particular interest in orbit closure containment and their singularities. See the surveys [Bon98, HZ14] for a detailed account. They are also important in Lie theory, where they lie at the foundation of Lusztig's geometric realization of Ringel's work on quantum groups [Lus90, Rin90]. From another viewpoint, quiver loci generalize some classically studied varieties such as determinantal varieties and varieties

---

2010 *Mathematics Subject Classification.* 14M12, 05E15, 14C17, 19E08.

Key words and phrases. quiver locus;representation variety;orbit closure;K-polynomial;multidegree;degeneracy locus;pipe dream;lacing diagram;matrix Schubert variety.

of complexes. This is because quiver loci are defined, at least up to radical, by minors of certain block form matrices built from the factors of $\mathtt{rep}_Q(\mathbf{d})$ (see the introduction and §3 of [KR15], or [RZ13]).

The line of approach most directly related to this paper was initiated by Buch and Fulton [BF99]. They produced formulas for equivariant cohomology classes of quiver loci, and interpreted them as universal formulas for degeneracy loci associated to representations of the quiver in the category of vector bundles on an algebraic variety. More formulas for equivariant cohomology and K-classes of quiver loci were subsequently produced in papers such as [Buc02, FR02, BFR05, KMS06]. Rimányi [Rim] has recently shown that these classes are natural structure constants for the Cohomological Hall Algebra of Kontsevich and Soibelman associated to the quiver [KS11]. A more detailed account of the state of the art can be found in recent works such as [Buc08, All14, Rim14].

1.2. **Summary of results and methods.** In this paper, we provide formulas for K-polynomials and multidegrees of quiver loci of type A quivers. There are three main steps in obtaining these formulas. The first step, which comprises the bulk of the paper, is to prove the various formulas in the bipartite (i.e. source-sink) orientation. The second step is to extend the bipartite results to all orientations using the connection between the arbitrarily oriented case and the bipartite case (see [KR15, §5]). To be precise, if Q is a type A quiver of arbitrary orientation, there is an associated bipartite type A quiver $\widetilde{Q}$ and a bijection between orbit closures for Q and a certain subset of orbit closures for $\widetilde{Q}$. We show that our formulas for the quiver loci of Q can be obtained by a simple substitution into the formulas for the corresponding quiver loci of $\widetilde{Q}$ (see Proposition 5.14). The third step is to simplify the various formulas to make them intrinsic to Q, so that the formulas make no reference to the added arrows in the associated bipartite quiver $\widetilde{Q}$ (see §5 for details). For the remainder of this subsection, we restrict to type A quivers of bipartite orientation, and discuss the formulas in this setting.

One thing that makes the bipartite orientation special is the bipartite Zelevinsky map constructed in [KR15]. It is an analogue of the map constructed by Zelevinsky for equioriented type A quivers in [Zel85], which was further studied in [LM98, KMS06]. In both cases, the maps give scheme theoretic isomorphisms between quiver loci and Kazhdan-Lusztig varieties, each of which is by definition the intersection of a Schubert variety and an opposite Schubert cell in a partial flag variety. The bipartite Zelevinsky map allows us to draw on the large body of knowledge about Schubert varieties to produce our formulas.

We now give a description of each formula and brief indication of the proof technique. In each case, we first prove the K-theoretic version. The associated multidegree version follows from the standard relation between K-polynomials and multidegrees (see §2.7). We state the formulas in the discussion here to give the reader an idea of their form, with the caveat that they depend on notations and conventions which will be established in later sections. At this point we just remark that each is modeled on the analogous formula for equioriented type A quivers in [KMS06]; a detailed literature comparison is found below in §1.3.

The *ratio formulas* are found in Theorem 3.1:

$$KQ_{\mathbf{r}}(\mathbf{t}/\mathbf{s}) = \frac{\mathfrak{G}_{v(\mathbf{r})}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})}{\mathfrak{G}_{v_*}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})} \qquad \text{and} \qquad \mathcal{Q}_{\mathbf{r}}(\mathbf{t}-\mathbf{s}) = \frac{\mathfrak{S}_{v(\mathbf{r})}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})}{\mathfrak{S}_{v_*}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})}.$$

They express each K-polynomial (resp., multidegree) as a ratio of double Grothendieck (resp., Schubert) polynomials. They are relatively straightforward consequences of the existence and properties of the bipartite Zelevinsky map.

The *pipe formulas* are found in Theorem 3.7:

$$K\mathcal{Q}_{\mathbf{r}}(\mathbf{t}/\mathbf{s}) = \sum_{P\in \mathrm{Pipes}(v_0,v(\mathbf{r}))} (-1)^{|P|-l(v(\mathbf{r}))}(\mathbf{1}-\mathbf{t}/\mathbf{s})^{P\setminus P_*}$$
$$\mathcal{Q}_{\mathbf{r}}(\mathbf{t}-\mathbf{s}) = \sum_{P\in \mathrm{RedPipes}(v_0,v(\mathbf{r}))} (\mathbf{t}-\mathbf{s})^{P\setminus P_*}.$$

They express each K-polynomial (resp., multidegree) as a sum over pipe dreams (resp., reduced pipe dreams) that have a certain shape related to the Zelevinsky map. Given the bipartite Zelevinsky map, their proofs also follow in a rather straightforward way from work of Woo and Yong on pipe formulas for Kazhdan-Lustzig varieties [WY12].

The *component formulas* are found in Theorem 4.37:

$$K\mathcal{Q}_{\mathbf{r}}(\mathbf{t}/\mathbf{s}) = \sum_{\mathbf{w}\in KW(\mathbf{r})} (-1)^{|\mathbf{w}|-\ell(v(\mathbf{r}))}\mathfrak{G}_{\mathbf{w}}(\mathbf{t};\mathbf{s})$$
$$\mathcal{Q}_{\mathbf{r}}(\mathbf{t}-\mathbf{s}) = \sum_{\mathbf{w}\in W(\mathbf{r})} \mathfrak{S}_{\mathbf{w}}(\mathbf{t};\mathbf{s}).$$

They express each K-polynomial (resp., multidegree) as a sum of products of Grothendieck (resp., Schubert) polynomials, where the sum is taken over K-theoretic (resp., minimal) lacing diagrams for the corresponding orbit closure. The proofs of the component formulas are more involved. The idea is to "degenerate" quiver loci to better understood varieties in a way that preserves K-polynomials and multidegrees. More precisely, we produce a flat family of group schemes acting fiberwise on a flat family of varieties such that over a general fiber, the orbit closures are isomorphic to the original quiver loci, and over the special fiber, the orbit closures are unions of products of matrix Schubert varieties. The multidegree formula can be obtained at this point, since it only requires knowledge of the irreducible components of the special fiber. The K-polynomial formula is more delicate: the coefficients of higher degree terms are determined by the Möbius function of the intersection poset of the irreducible components.

Remark 1.1. Although we use the algebraic language of multidegrees and K-polynomials in this paper, our formulas also hold in other settings since there are several other interpretations of the multidegrees and K-polynomials that we study. These alternate interpretations are in the languages of equivariant cohomology and K-theory, [KMS06, §1.4], and the virtual, rational representations of GL [Buc08, §3]. When cited literature is written from one of these perspectives, we will use the K-polynomial or multidegree version without explicit mention of the conversion. □

1.3. **Relation to existing literature.** Most of the formulas for quiver loci in the literature are for the equioriented, type A case. As mentioned above, our formulas and proof techniques are directly modeled on [KMS06]. There one already finds a K-theoretic ratio formula for the equioriented case; we have generalized this to arbitrary orientation. Our pipe formulas and component formulas generalize theirs in two directions: by moving from a specific orientation to arbitrary orientation, and also by moving from multidegrees to K-polynomials. Some intermediate results that helped us find these generalizations are the K-theoretic pipe formula [Mil05, Theorem 3] and component formula [Buc05, Theorem 6.3] for the equioriented case, and the multidegree component formula for arbitrary orientation [BR07, Theorem 1]. The K-theoretic component formula that we prove in §5 is Conjecture 1 of [BR07].

Our proofs of the component formulas in the bipartite setting use the Gröbner degeneration ideas of [KMS06] (cf. the later paper [Yon05] for a purely combinatorial approach), however, our proof is more direct by taking advantage of recent works [WY12, Knu]. The proof of the equioriented type A component formula given in [KMS06] involves taking a certain limit as the dimension vector grows, obtained by adding copies of the projective-injective indecomposable representation of an equioriented type A quiver (the "longest lace"). Other orientations never have a projective-injective representation, and it is not clear to us what the analogous technique would be. In particular, this is why we do not give an analogue of the fourth positive formula from [KMS06], the tableau formula. That formula is given in terms of Schur functions, and its proof depends on the stable component formula in terms of Stanley symmetric functions. This formula is most closely related to the conjectured positive formula for quiver loci of all Dynkin types in [Buc08, Conjecture 1.1].

Finally, we note that our work gives a geometric interpretation of the "double" quiver polynomials studied in [Buc02, KMS06] and other works cited above: these double quiver polynomials come from the natural action of a larger torus on the equioriented orbit closures, after embedding them into a larger representation space associated to a bipartite quiver.

Acknowledgements. We would like to thank Alex Yong for directing us to his work with Alex Woo, and Christian Stump for his pipe dream LaTeX macros. We also thank Anders Buch and Alex Yong for their comments on a draft of this paper, and the anonymous referees for numerous helpful comments and suggestions.

2. BACKGROUND AND PRELIMINARY RESULTS

We work over a fixed field K throughout the paper, which is often omitted from our notation. To simplify the exposition below, we will assume that K is algebraically closed. However, since all schemes appearing in this paper are defined over $\mathbb{Z}$, there is no difficulty generalizing results to arbitrary K.

2.1. Type A quiver loci.

Fix a quiver $Q = (Q_0, Q_1)$ with vertex set Q_0 and arrow set Q_1. Given a dimension vector $\mathbf{d}\colon Q_0 \to \mathbb{Z}_{\geq 0}$, we have the *representation space*

$$\mathtt{rep}_Q(\mathbf{d}) := \prod_{a \in Q_1} \mathrm{Mat}(\mathbf{d}(ta), \mathbf{d}(ha)), \tag{2.1}$$

where $\mathrm{Mat}(m, n)$ denotes the algebraic variety of matrices with m rows, n columns, and entries in K, and ta and ha denote the tail and head of an arrow $ta \xrightarrow{a} ha$. Each $V = (V_a)_{a\in Q_1}$ in $\mathtt{rep}_Q(\mathbf{d})$ is a *representation* of Q; each matrix V_a maps row vectors in $K^{\mathbf{d}(ta)}$ to row vectors in $K^{\mathbf{d}(ha)}$ by right multiplication. An introduction to the theory of quiver representations can be found in the textbooks [ASS06, Sch14]. Observe that $\mathtt{rep}_Q(\mathbf{d})$ parametrizes all representations of Q with vector space $K^{\mathbf{d}(z)}$ at vertex $z \in Q_0$. We write the corresponding total dimension as $d = \sum_{z\in Q_0} \mathbf{d}(z)$.

There is a *base change group*

$$GL(\mathbf{d}) := \prod_{z\in Q_0} GL_{\mathbf{d}(z)}, \tag{2.2}$$

which acts on $\mathtt{rep}_Q(\mathbf{d})$ by change of basis. Here $GL_{\mathbf{d}(z)}$ denotes the general linear group of $\mathbf{d}(z) \times \mathbf{d}(z)$ matrices with entries in K. Explicitly, if $g = (g_z)_{z\in Q_0}$ is an element of $GL(\mathbf{d})$, and $V = (V_a)_{a\in Q_1}$ is an element of $\mathtt{rep}_Q(\mathbf{d})$, then the (right) action of $GL(\mathbf{d})$ on $\mathtt{rep}_Q(\mathbf{d})$ is given by

$$V \cdot g = (g_{ta}^{-1} V_a g_{ha})_{a\in Q_1}. \tag{2.3}$$

Two points $V, W \in \mathtt{rep}_Q(\mathbf{d})$ lie in the same orbit if and only if V and W are isomorphic as representations of Q. The closure of a $GL(\mathbf{d})$-orbit in $\mathtt{rep}_Q(\mathbf{d})$ is called a *quiver locus*.

In this paper, we only work with quivers of Dynkin type A. We arbitrarily designate one endpoint “left” and the other “right” so that we can speak of arrows pointing left or right. The fundamental orientation in type A, to which understanding the geometry of quiver loci in all other orientations essentially reduces, is the *bipartite orientation*, where every vertex is either a sink or source (see [KR15, §5] for details). In this setting, we label all sink vertices by x_j, all source vertices by y_i, leftward pointing arrows by α_j, and rightward pointing arrows by β_i. Since we act on row vectors instead of column vectors in this paper, this notation slightly differs from [KR15]. We use the following running example throughout the paper.

$$Q = \begin{tikzcd} y_3 \arrow[dr, "\beta_3"'] & & y_2 \arrow[dl, "\alpha_3"'] \arrow[dr, "\beta_2"'] & & y_1 \arrow[dl, "\alpha_2"'] \arrow[dr, "\beta_1"'] & & y_0 \arrow[dl, "\alpha_1"'] \\ & x_3 & & x_2 & & x_1 & \end{tikzcd} \tag{2.4}$$

We will assume that all of our bipartite type A quivers start and end with a source vertex, as this will simplify the exposition. This is harmless, because one can always extend any quiver on either end by a vertex z of dimension 0 (i.e. $\mathbf{d}(z) = 0$).

The orbits in $\mathtt{rep}_Q(\mathbf{d})$ (and their closures) are indexed by *rank arrays* [ADF85] (see also [KR15, §3.2]). These collections of integers encode dimensions of Hom spaces between representations. We now recall the relevant details in the bipartite setting.

Assume Q is a bipartite type A quiver with n leftward pointing arrows α_i, and n rightward pointing arrows β_j. Given a representation $V \in \mathtt{rep}_Q(\mathbf{d})$, define $M(V)$ to be the matrix

$$M(V) = \begin{bmatrix} & & & & V_{\alpha_1} \\ & & & V_{\alpha_2} & V_{\beta_1} \\ & & V_{\alpha_3} & V_{\beta_2} & \\ & \iddots & \iddots & & \\ V_{\alpha_n} & V_{\beta_{n-1}} & & & \\ V_{\beta_n} & & & & \end{bmatrix}, \tag{2.5}$$

with 0 entries in the unlabeled blocks.

Given an interval $J = [\gamma, \delta] \subseteq Q$ having leftmost arrow γ and rightmost arrow δ, let $M_J(V)$ be the submatrix of $M(V)$ whose upper right block is associated to δ and whose lower left block is associated to γ. For example, we have

$$M_{[\alpha_5,\alpha_3]}(V) = \begin{bmatrix} & & V_{\alpha_3} \\ & V_{\alpha_4} & V_{\beta_3} \\ V_{\alpha_5} & V_{\beta_4} & \end{bmatrix} \quad \text{and} \quad M_{[\alpha_4,\beta_2]}(V) = \begin{bmatrix} & V_{\alpha_3} & V_{\beta_2} \\ V_{\alpha_4} & V_{\beta_3} & \end{bmatrix}. \tag{2.6}$$

The rank of such a matrix is invariant under the action of the base change group, so each interval J defines a function which is constant on orbits.

$$\mathbf{r}_J \colon \mathtt{rep}_Q(\mathbf{d}) \to \mathbb{Z}_{\geq 0}, \quad r_J(V) = \operatorname{rank} M_J(V) \tag{2.7}$$

In fact, two representations $V, W \in \mathtt{rep}_Q(\mathbf{d})$ lie in the same $GL(\mathbf{d})$-orbit if and only if $\mathbf{r}_J(V) = \mathbf{r}_J(W)$ for all intervals $J \subseteq Q$. Define the *rank array of the representation* V (or, equivalently, of the $GL(\mathbf{d})$-orbit through V) to be the collection $\mathbf{r} = (\mathbf{r}_J(V))_{J \subseteq Q}$. Throughout this paper, we use the notation $\Omega_{\mathbf{r}}$ to denote the closure of the $GL(\mathbf{d})$-orbit with rank array $\mathbf{r}$.

Remark on notation: Until §5, we work with one fixed bipartite type A quiver Q and dimension vector $\mathbf{d}$ at a time; hence, these will be omitted from notation whenever possible. In particular, we write $\mathtt{rep}$ instead of $\mathtt{rep}_Q(\mathbf{d})$, and GL instead of $GL(\mathbf{d})$.

2.2. **Permutations and permutation matrices.** We now fix our conventions regarding permutations and permutation matrices. A permutation v in the symmetric group S_n is a bijection $v\colon [1,n] \to [1,n]$. If $v, w \in S_n$, then $vw \in S_n$ will denote the composition of functions $i \mapsto v(w(i))$, $1 \leq i \leq n$. A *simple transposition* s_i is a permutation that switches i and $i+1$, while fixing everything else, and the *length* $\ell(v)$ of a permutation v is the minimal number of factors needed to express v as a product of simple transpositions.

To the permutation v, one can associate an $n \times n$ permutation matrix v^T which has a 1 in location (i,j) if $v(i) = j$, and a 0 in location (i,j) otherwise. Using this convention, it is easy to check that

$$\vec{e}_i v^T = \vec{e}_{v(i)}$$

where $\vec{e}_i$ is the $1 \times n$ row vector with a 1 in position i and 0s elsewhere.

To simplify the notation in the paper, we will often use v to denote a both a permutation and its associated matrix. It will be clear from the context whether we are working with a function or a matrix.

2.3. **The bipartite Zelevinsky map.** In this section, we recall the relationship between quiver loci of bipartite type A quivers and Schubert varieties which was established in [KR15]. Fix a bipartite type A quiver Q with $n+1$ source vertices $y_0, y_1, \ldots, y_n$, labelled from right to left, and n sink vertices $x_1, \ldots, x_n$, labelled from right to left (e.g., see (2.4) above). Fix a dimension vector $\mathbf{d}$, retaining the notation of §2.1. Let

$$d_x = \sum_{j=1}^{n} \mathbf{d}(x_j) \qquad\qquad d_y = \sum_{i=0}^{n} \mathbf{d}(y_i), \tag{2.8}$$

so $d = d_x + d_y$. Let $Y_\circ^{v_0}$ denote the space of $d \times d$ matrices of the form

$$\begin{bmatrix} * & \mathbf{1}_{d_y} \\ \mathbf{1}_{d_x} & 0 \end{bmatrix} \tag{2.9}$$

where the entries in the $*$ block are arbitrary elements of the base field K.

Let Z be a matrix of the form in (2.9), which has indeterminates (instead of field elements) as entries in the $*$ block. We may thus identify the coordinate ring $K[Y_\circ^{v_0}]$ with the polynomial ring $K[z_{ij}] := K[z_{ij} : 1 \leq i \leq d_y,\ 1 \leq j \leq d_x]$ generated by the indeterminates in Z. Let $Z_{p\times q}$ denote the northwest submatrix of Z consisting of the top p rows and left q columns (*not* necessarily inside the $*$ block). Given a $d \times d$ permutation matrix v, define the ideal $I_v \subseteq K[z_{ij}]$ by

$$I_v := \langle \text{minors of size } (1 + \operatorname{rank} v_{p\times q}) \text{ in } Z_{p\times q} \mid (p,q) \in [1,d] \times [1,d] \rangle.$$

The affine scheme defined by the prime ideal I_v, which we denote by $Y_v \subseteq Y_\circ^{v_0}$, is called a *Kazhdan-Lusztig variety* and is isomorphic to the intersection of an opposite Schubert cell and a Schubert variety (see [WY08] for details). To be precise, let $G := GL_d$ and let P be the parabolic subgroup of block lower triangular matrices where the diagonals have block sizes

$$\mathbf{d}(y_0), \ldots, \mathbf{d}(y_n),\ \mathbf{d}(x_n),\ \mathbf{d}(x_{n-1}), \ldots, \mathbf{d}(x_1), \tag{2.10}$$

listed from northwest to southeast. Then $Y_\circ^{v_0}$ is isomorphic to the opposite Schubert cell $P\backslash Pv_0B^-$, where B^- is the Borel subgroup of lower triangular matrices, and v_0 is the permutation matrix

$$v_0 := \begin{bmatrix} 0 & \mathbf{1}_{d_y} \\ \mathbf{1}_{d_x} & 0 \end{bmatrix}. \tag{2.11}$$

The affine scheme Y_v is isomorphic to the intersection of the Schubert variety $\overline{P\backslash PvB^+}$ with the opposite cell $P\backslash Pv_0B^-$.

$$\zeta(V) = \begin{array}{c|ccc|cccc|}
 & \mathbf{d}(x_3) & \mathbf{d}(x_2) & \mathbf{d}(x_1) & \mathbf{d}(y_0) & \mathbf{d}(y_1) & \mathbf{d}(y_2) & \mathbf{d}(y_3) \\
\mathbf{d}(y_0) & 0 & 0 & V_{\alpha_1} & \mathbf{1}_{\mathbf{d}(y_0)} & & & \\
\mathbf{d}(y_1) & 0 & V_{\alpha_2} & V_{\beta_1} & & \mathbf{1}_{\mathbf{d}(y_1)} & & \\
\mathbf{d}(y_2) & V_{\alpha_3} & V_{\beta_2} & 0 & & & \mathbf{1}_{\mathbf{d}(y_2)} & \\
\mathbf{d}(y_3) & V_{\beta_3} & 0 & 0 & & & & \mathbf{1}_{\mathbf{d}(y_3)} \\
\cline{2-8}
\mathbf{d}(x_3) & \mathbf{1}_{\mathbf{d}(x_3)} & & & & & & \\
\mathbf{d}(x_2) & & \mathbf{1}_{\mathbf{d}(x_2)} & & & & \mathbf{0} & \\
\mathbf{d}(x_1) & & & \mathbf{1}_{\mathbf{d}(x_1)} & & & &
\end{array}$$

FIGURE 1. Image of the Zelevinsky map in $Y_\circ^{v_0}$, labeled by block sizes

In [KR15], it is shown that every bipartite type A quiver locus is isomorphic to a Kazhdan-Lusztig variety $Y_v \subseteq Y_\circ^{v_0}$. This is proven by defining a closed immersion $\zeta\colon \texttt{rep} \to Y_\circ^{v_0}$, which we call a *bipartite Zelevinsky map*. The image of a representation $(V_a)_{a \in Q_1}$ under this map is shown in Figure 1 in the case of our running example. The definition of the bipartite Zelevinsky map for an arbitrary bipartite type A quiver is extrapolated in the obvious way. In particular, if $V \in \texttt{rep}$, and $M(V)$ is as in (2.5), then

$$\zeta(V) = \begin{bmatrix} M(V) & \mathbf{1}_{d_y} \\ \mathbf{1}_{d_x} & 0 \end{bmatrix}$$

We then have:

Theorem 2.12. [KR15, Theorem 4.12] *The bipartite Zelevinsky map ζ restricts to a scheme-theoretic isomorphism from each orbit closure $\Omega_{\mathbf{r}} \subseteq$ $\texttt{rep}$ to a Kazhdan-Lusztig variety $Y_{v(\mathbf{r})} \subseteq Y_\circ^{v_0}$.*

The permutation $v(\mathbf{r})$ is known as the *Zelevinsky permutation* of the quiver locus $\Omega_{\mathbf{r}}$; its definition is given in §2.4 below and an equivalent characterization given in Theorem 2.33. We define v_* to be the Zelevinsky permutation of the entire space $\texttt{rep}$, so Y_{v_*} is exactly the image of ζ.

Definition 2.13. The blocks in the northwest quadrant of a generic matrix in Y_{v_*} which are not identically zero are referred to as the *snake region*. □

For example, in Figure 1, the blocks labelled by V_{α_i} or V_{β_j} are the blocks in the snake region.

2.4. **Zelevinsky permutations.**

In this section, we recall the definition of the *bipartite Zelevinsky permutation* $v(\mathbf{r})$ from [KR15, §4], which determines the Kazhdan-Lusztig variety $Y_{v(\mathbf{r})}$ in Theorem 2.12 above. This definition of $v(\mathbf{r})$ goes through an intermediate concept called a *block rank matrix*, which we recall here to make the paper more self contained, although it is not used elsewhere in this work. We give an equivalent characterization of the Zelevinsky permutation in terms of lacing diagrams in §2.10 below, which is more closely related to the combinatorics of this paper. We retain the notation of the previous section.

A dimension vector $\mathbf{d}$ determines a subdivision of any $d \times d$ matrix into blocks. Row blocks are indexed from top to bottom by $y_0, \ldots, y_n, x_n, x_{n-1}, \ldots, x_1$, with sizes determined by $\mathbf{d}$ (i.e., as in (2.10) above). Column blocks are indexed from left to right by $x_n, x_{n-1}, \ldots, x_1, y_0, y_1, \ldots, y_n$, with sizes similarly determined by $\mathbf{d}$ (see Figure 1)

$$\mathbf{d}(x_n), \ldots, \mathbf{d}(x_1),\ \mathbf{d}(y_0),\ \mathbf{d}(y_1), \ldots, \mathbf{d}(y_n). \tag{2.14}$$

For such a matrix M, let $M_{i\times j}$ denote the submatrix of consisting of the topmost i block rows, and leftmost j block columns.

Let $\mathbf{r}$ be a quiver rank array and $V \in \texttt{rep}(\mathbf{d})$ a representation in the orbit associated to $\mathbf{r}$. The *northwest block rank matrix* $\mathbf{b}(\mathbf{r})$ is the $(2n+1)\times(2n+1)$ matrix $\mathbf{b}(\mathbf{r})$ defined by

$$\mathbf{b}(\mathbf{r})_{i,j} = \operatorname{rank}\left(\zeta(V)_{i\times j}\right)$$

where $\zeta(V)$ denotes the Zelevinsky image of V in $Y_\circ^{v_0}$. This is well-defined; any representation with rank array $\mathbf{r}$ gives rise to the same block rank matrix $\mathbf{b}(\mathbf{r})$. Furthermore, the combinatorial data in a quiver rank array $\mathbf{r}$ and northwest block rank matrix $\mathbf{b}(\mathbf{r})$ are equivalent, in the sense that a matrix $M \in Y_\circ^{v_0}$ satisfies

$$\operatorname{rank} M_{i\times j} = \mathbf{b}(\mathbf{r})_{i,j} \quad \text{for all } i,j$$

if and only if $M = \zeta(V)$ for some V with rank array $\mathbf{r}$.

The *Zelevinsky permutation* $v(\mathbf{r})$ *associated to rank array* $\mathbf{r}$ is then defined to be the permutation in S_d which satisfies the following conditions:

(1) the number of 1s in block (i,j) of the $d\times d$ permutation matrix $v(\mathbf{r})^T$ is equal to

$$\mathbf{b}(\mathbf{r})_{i,j} + \mathbf{b}(\mathbf{r})_{i-1,j-1} - \mathbf{b}(\mathbf{r})_{i,j-1} - \mathbf{b}(\mathbf{r})_{i-1,j}$$

where $\mathbf{b}(\mathbf{r})_{i,j} = 0$ if i or j is outside of the range $[1, 2n+1]$;

(2) the 1s in $v(\mathbf{r})^T$ are arranged from northwest to southeast across each block row;

(3) the 1s in $v(\mathbf{r})^T$ are arranged from northwest to southeast down each block column.

These conditions uniquely determine $v(\mathbf{r})$ [KR15, Prop. 4.8].

Example 2.15. Continuing our running example, we further take $\mathbf{d} = (2,2,2,3,2,2,1)$. Consider the representation $V = (V_{\beta_3}, V_{\alpha_3}, V_{\beta_2}, V_{\alpha_2}, V_{\beta_1}, V_{\alpha_1})$

(2.16)
$$\begin{tikzcd}
K^2 \arrow[dr, swap, "V_{\beta_3}"] & & K^2 \arrow[dl, swap, "V_{\alpha_3}"] \arrow[dr, "V_{\beta_2}"] & & K^2 \arrow[dl, swap, "V_{\alpha_2}"] \arrow[dr, "V_{\beta_1}"] & & K \arrow[dl, "V_{\alpha_1}"] \\
& K^2 & & K^3 & & K^2 &
\end{tikzcd}$$

given by matrices

(2.17)
$$V = \left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \end{bmatrix} \right).$$

Then we can directly compute the block rank matrix

(2.18)
$$\mathbf{b}(\mathbf{r}) = \left[\begin{array}{ccc|cccc}
0 & 0 & 1 & 1 & 1 & 1 & 1 \\
0 & 2 & 3 & 3 & 3 & 3 & 3 \\
2 & 4 & 5 & 5 & 5 & 5 & 5 \\
2 & 4 & 5 & 6 & 6 & 6 & 7 \\
\hline
2 & 4 & 6 & 7 & 7 & 7 & 9 \\
2 & 5 & 7 & 8 & 8 & 10 & 12 \\
2 & 5 & 7 & 8 & 10 & 12 & 14
\end{array}\right].$$

One constructs $v(\mathbf{r})^T$ from $\mathbf{b}(\mathbf{r})$ by working from the northwest to southeast, filling in each block with the number of 1s specified in item (1) above, arranged according to (2) and (3).

The empty blocks in the matrix below contain all zeros.

$$v(\mathbf{r})^T = \left[\begin{array}{cc|ccc|cc|c|cc|cc|cc}
 & & & & & 1 & 0 & & & & & & & \\
\hline
 & & 1 & 0 & 0 & & & & & & & & & \\
 & & 0 & 1 & 0 & & & & & & & & & \\
\hline
1 & 0 & & & & & & & & & & & & \\
0 & 1 & & & & & & & & & & & & \\
\hline
 & & & & & & & 1 & & & & & 0 & 0 \\
 & & & & & & & 0 & & & & & 1 & 0 \\
\hline
 & & & & & 0 & 1 & & & & & & 0 & 0 \\
 & & & & & 0 & 0 & & & & & & 0 & 1 \\
\hline
 & & 0 & 0 & 1 & & & & & & 0 & 0 & & \\
 & & 0 & 0 & 0 & & & & & & 1 & 0 & & \\
 & & 0 & 0 & 0 & & & & & & 0 & 1 & & \\
\hline
 & & & & & & & & 1 & 0 & & & & \\
 & & & & & & & & 0 & 1 & & & &
\end{array}\right] \tag{2.19}$$

□

2.5. **Matrix Schubert varieties.** In our discussion of the component formulas, we encounter matrix Schubert varieties [Ful92]. These are subvarieties of the space of $k \times l$ matrices which are obtained by imposing rank conditions on certain submatrices.

Let $X = \mathrm{Mat}(k, l)$, the space of $k \times l$ matrices with entries in the field K, and let v be a partial permutation matrix in X (i.e. a $k \times l$ matrix of 1s and 0s with at most one 1 in each row and column). Given $M \in X$, let $M_{p\times q}$ denote the submatrix of M consisting of the intersection of the top p rows and left q columns. The *northwest matrix Schubert variety* X_v is the subvariety of X defined as

$$X_v := \{M \in X \mid \mathrm{rank}\ M_{p\times q} \leq \mathrm{rank}\ v_{p\times q}\}.$$

Let $M^{p\times q}$ denote the submatrix of $M \in X$ consisting of the intersection of the bottom p rows and right q columns. The *southeast matrix Schubert variety* X^v is the subvariety of X defined as

$$X^v := \{M \in X \mid \mathrm{rank}\ M^{p\times q} \leq \mathrm{rank}\ v^{p\times q}\}.$$

If v_{rot} denotes the 180° rotation of the matrix v, then $X^v \cong X_{v_{\mathrm{rot}}}$.

Let $Z = (z_{ij})$ be a $k \times l$ matrix of indeterminates. The (prime) defining ideals of X_v and X^v are

$$I_v := \langle \text{minors of size } (1 + \mathrm{rank}\, v_{p\times q}) \text{ in } Z_{p\times q} \mid (p, q) \in [1, k] \times [1, l]\rangle$$

and

$$I^v := \langle \text{minors of size } (1 + \mathrm{rank}\, v^{p\times q}) \text{ in } Z^{p\times q} \mid (p, q) \in [1, k] \times [1, l]\rangle$$

respectively.

We now describe a smaller set of generators for the ideals I_v and I^v. Let v be a $k \times l$ partial permutation matrix, and assign to it a $k \times l$ grid. The set of locations (or boxes) in the grid that have a 1 neither directly north nor directly west is called the *(Rothe) diagram* of v, denoted $\mathcal{D}_v$. Fulton's *essential set* $\mathcal{E}ss(v)$ is the set of those $(i, j) \in \mathcal{D}_v$ such that neither $(i + 1, j)$ nor $(i, j + 1)$ is also in the diagram of v. By [Ful92, §3], we have

$$I_v = \langle \text{minors of size } (1 + \mathrm{rank}\, v_{p\times q}) \text{ in } Z_{p\times q} \mid (p, q) \in \mathcal{E}ss(v)\rangle.$$

Analogously, one can check that

$$I^v = \langle \text{minors of size } (1 + \mathrm{rank}\, v^{p\times q}) \text{ in } Z^{p\times q} \mid (k - p + 1, l - q + 1) \in \mathcal{E}ss(v_{\mathrm{rot}})\rangle.$$

Each variety X_v and X^v is an orbit closure for the action of a product of Borel subgroups: let B_+ and B_- denote Borel subgroups of invertible upper and lower triangular matrices respectively. Then we have that

$$X_v = \overline{B_- v B_+} \text{ and } X^v = \overline{B_+ v B_-},$$

where the closures are taken inside of the space X of $k \times l$ matrices, the Borel subgroups to the left of v are inside of GL_k, and the Borel subgroups to the right of v are inside of GL_l.

Let v be a $k \times l$ partial permutation matrix. Define the *completion* $c(v)$ of v to be the permutation matrix which is (i) of minimal possible size such that v lies in the northwest corner (i.e. $c(v)_{k\times l} = v$), and (ii) has minimal length among such permutation matrices. Define the *opposite completion* $oc(v)$ of v to be the permutation matrix which is (i) of minimal possible size such that v lies in the southeast corner (i.e. $oc(v)^{k\times l} = v$), and (ii) has minimal length among such permutation matrices. In other words, the opposite completion of v is obtained by rotating v by 180°, completing to a permutation matrix, and then rotating back by 180°.

Finally, it will be useful to note that (i,j) is in the essential set of a partial permutation v if and only (i,j) is in the essential set of the completion of v. Consequently, the ideals I_v and $I_{c(v)}$ have a generating set in common (namely, the one coming from Fulton's essential minors), although the latter ideal is in a polynomial ring in more variables. Analogously, I^v and $I^{oc(v)}$ have a generating set in common.

2.6. **Multigradings.**

There are natural torus actions on the varieties central to this paper, and these torus actions induce multigradings on their coordinate rings in the standard way. We begin this subsection by a brief explanation of this translation.

Suppose we have an algebraic left action of an algebraic torus $T \simeq (K^\times)^d$ on an affine variety $X = \operatorname{Spec} R$, with the action of $t \in T$ on $x \in X$ written as $t \cdot x$. This induces a left action of T on the K-algebra R, with the correspondence given by $(t\cdot f)(x) = f(t^{-1}\cdot x)$ for $f \in R$. For each character $\varphi\colon T \to K^\times$, we have a *weight space*

$$R_\varphi = \{f \in R \mid t \cdot f = \varphi(t) f\}, \tag{2.20}$$

and a decomposition of R as a K-vector space

$$R \simeq \bigoplus_{\varphi} R_\varphi. \tag{2.21}$$

It is straightforward to verify that this decomposition is actually a grading of R by the character group of T. If we take an isomorphism $T \simeq (K^\times)^d$, writing $t = (t_1, \dots, t_d)$ with each $t_i \in K^\times$, this identifies the character group of T with $\mathbb{Z}^d$: each character φ is of the form $\varphi(t) = t_1^{e_1}\cdots t_d^{e_d}$ and thus φ is identified with $(e_1, \dots, e_d) \in \mathbb{Z}^d$, and so the torus action gives a $\mathbb{Z}^d$-multigrading to R. Working in reverse, a $\mathbb{Z}^d$-multigrading of an affine K-algebra R determines an algebraic action of the torus $(K^\times)^d$ on the variety $\operatorname{Spec} R$. A closed subvariety $\operatorname{Spec}(R/I) \subseteq X$ is T-stable exactly when I is homogeneous with respect to this multigrading. Given the equivalence of these concepts, we may sometimes only explicitly describe one or the other, but use both, in various parts of the paper.

Let Q be any quiver and let $\mathbf{d}$ be a dimension vector for Q. Let T be the maximal torus of $GL(\mathbf{d})$ consisting of matrices which are diagonal in each factor. The restriction of the $GL(\mathbf{d})$-action on $\mathtt{rep}_Q(\mathbf{d})$ to T induces a multigrading on $K[\mathtt{rep}_Q(\mathbf{d})]$ as in the previous paragraph, which makes the ideals of orbit closures homogeneous since orbit closures are T-stable. Now we introduce notation to explicitly describe this for bipartite quivers, deferring the case of arbitrary orientation until §5.

$$\begin{array}{c|ccc|cccc|}
 & \mathbf{s}^3 & \mathbf{s}^2 & \mathbf{s}^1 & \mathbf{t}^0 & \mathbf{t}^1 & \mathbf{t}^2 & \mathbf{t}^3 \\
\mathbf{t}^0 & * & * & * & \mathbf{1}_{\mathbf{d}(y_0)} & & & \\
\mathbf{t}^1 & * & * & * & & \mathbf{1}_{\mathbf{d}(y_1)} & & \\
\mathbf{t}^2 & * & * & * & & & \mathbf{1}_{\mathbf{d}(y_2)} & \\
\mathbf{t}^3 & * & * & * & & & & \mathbf{1}_{\mathbf{d}(y_3)} \\
\hline
\mathbf{s}^3 & \mathbf{1}_{\mathbf{d}(x_3)} & & & & & & \\
\mathbf{s}^2 & & \mathbf{1}_{\mathbf{d}(x_2)} & & & & \mathbf{0} & \\
\mathbf{s}^1 & & & \mathbf{1}_{\mathbf{d}(x_1)} & & & &
\end{array}$$

FIGURE 2. Alphabets assigned to each block in multigrading of $Y_\circ^{v_0}$

To explicitly describe the multigrading of $K[\mathtt{rep}]$ in the bipartite orientation, we associate an alphabet $\mathbf{s}^j$ to the vertex x_j, and an alphabet $\mathbf{t}^i$ to the vertex y_i:

$$\mathbf{s}^j = s_1^j, s_2^j, \ldots, s_{\mathbf{d}(x_j)}^j \quad \text{and} \quad \mathbf{t}^i = t_1^i, t_2^i, \ldots, t_{\mathbf{d}(y_i)}^i. \tag{2.22}$$

Let $\mathbf{t}$ be the sequence $\mathbf{t}^0, \mathbf{t}^1, \ldots, \mathbf{t}^n$ and $\mathbf{s}$ be the sequence $\mathbf{s}^n, \ldots, \mathbf{s}^2, \mathbf{s}^1$ (the order in which the alphabets are concatenated is indicated, as in Figure 2, with each individual alphabet still in its standard order). Recall d_x and d_y from (2.8) and that $d = d_x + d_y$. Identify $\mathbb{Z}^{d_x}$ with the free abelian group on $\mathbf{s}$ and $\mathbb{Z}^{d_y}$ with the free abelian group on $\mathbf{t}$, and $\mathbb{Z}^d = \mathbb{Z}^{d_x} \oplus \mathbb{Z}^{d_y}$. Then the induced multigrading of the coordinate ring $K[\mathtt{rep}]$ by $\mathbb{Z}^d$ has the coordinate function $f_{ij}^{\alpha_k}$, which picks out the entry in row i and column j of V_{α_k}, in degree $t_i^{k-1} - s_j^k$. Similarly, the coordinate function $f_{ij}^{\beta_k}$ picking out the entry in row i and column j of V_{β_k} has degree $t_i^k - s_j^k$.

Our multigrading of $K[Y_\circ^{v_0}]$ by $\mathbb{Z}^d$ is induced by the right action of the torus of diagonal matrices $T \subseteq GL_d$ on $P \backslash P v_0 B^-$. To explicitly describe it, recall the block structure for matrices in $Y_\circ^{v_0}$ introduced in §2.4. Consider the submatrix of indeterminates in the block with row index y_k and column index x_l. Then, if h is the coordinate function picking out the entry in the (i,j)-location of this submatrix, give h degree $t_i^k - s_j^l$. Figure 2 illustrates a labeling of row and column blocks of $Y_\circ^{v_0}$ by the alphabets from (2.22). The degree of a coordinate function h as above is its row label minus its column label.

Lemma 2.23. *The bipartite Zelevinsky map ζ of §2.3 is $GL(\mathbf{d})$-equivariant with respect to the embedding $GL(\mathbf{d}) \subset GL_d$ as block diagonal matrices with blocks of sizes and order given in (2.14). In particular, it is T-equivariant also, so for each rank array $\mathbf{r}$, the induced isomorphism of coordinate rings $\zeta^\sharp : K[\zeta(\Omega_\mathbf{r})] \to K[\Omega_\mathbf{r}]$ is multigraded.*

Proof. The isomorphism $Y_\circ^{v_0} \simeq P \backslash P v_0 B^-$ is essentially a choice of P-coset representatives, which is not stable under right multiplication by $GL(\mathbf{d})$. So when $GL(\mathbf{d})$ acts on an element of $Y_\circ^{v_0}$ by right multiplication, the left P-action is necessary to get a representative in $Y_\circ^{v_0}$. The situation is illustrated in (2.24), where $\sim$ indicates equivalence modulo P (which acts by row operations).

$$\begin{bmatrix} M & \mathbf{1}_{d_y} \\ \mathbf{1}_{d_x} & 0 \end{bmatrix} \begin{bmatrix} g_x & 0 \\ 0 & g_y \end{bmatrix} = \begin{bmatrix} M g_x & g_y \\ g_x & 0 \end{bmatrix} \sim \begin{bmatrix} g_y^{-1} M g_x & \mathbf{1}_{d_y} \\ \mathbf{1}_{d_x} & 0 \end{bmatrix} \tag{2.24}$$

For $g = diag(g_x, g_y) \in GL(\mathbf{d})$ and $V \in \texttt{rep}$, the expression (2.3) for $g \cdot V$ shows that the block form of g_x and g_y will make $g_y^{-1} M g_x$ exactly the northwest quadrant of $\zeta(g \cdot V)$. So ζ is $GL(\mathbf{d})$-equivariant. □

Let $X = \mathrm{Mat}(k, l)$. A matrix Schubert variety $X_v \subseteq X$ has an action of the diagonal torus $(K^\times)^k \times (K^\times)^l \subset GL_k \times GL_l$. For an explicit description of the multigrading of $K[X_v]$, assign letters $\mathbf{a} = a_1, \dots, a_k$ to the rows, from top to bottom, and letters $\mathbf{b} = b_1, \dots, b_l$ to the columns, from left to right. The coordinate function picking out the (i, j) entry of a matrix in X_v will have degree $a_i - b_j$. For opposite matrix Schubert varieties, the same torus acts, but the isomorphism $X^v \cong X_{v_{\mathrm{rot}}}$ is not T-equivariant with respect to this action because an automorphism of T is needed. We will use the symbol $\widetilde{\ }$ over an alphabet to denote that the order is reversed; with this notation, an isomorphism $X^v \cong X_{v_{\mathrm{rot}}}$ becomes multigraded when accompanied by substituting $\widetilde{\mathbf{a}}$ for $\mathbf{a}$ and $\widetilde{\mathbf{b}}$ for $\mathbf{b}$.

2.7. **K-polynomials and multidegrees.** Here we give a brief introduction to K-polynomials, following [MS05, Chapter 8]. Geometrically, these polynomials represent classes in equivariant K-theory. Let S be a polynomial ring over K, multigraded by $\mathbb{Z}^d$. Assume the multigrading is *positive* (which is the case for all multigradings in this paper), meaning that the only elements of degree $\mathbf{0}$ are the constants in the base field K. Let $\mathbf{a} = a_1, \dots, a_d$ be an alphabet with d letters, identifying $\mathbb{Z}^d$ with the free abelian group on $\mathbf{a}$. For each $\mathbf{e} = \sum e_i a_i \in \mathbb{Z}^d$, where $e_i \in \mathbb{Z}$, we define the monomial $\mathbf{a}^{\mathbf{e}} = a_1^{e_1} \cdots a_d^{e_d}$. Note that $\mathbf{e}$ can be recovered from $\mathbf{a}^{\mathbf{e}}$ by substituting $1 - a_i$ for each a_i, substituting the power series expansion $(1 - a_i)^{-1} = \sum_{j \geq 0} a_i^j$ as necessary to get all positive exponents, and taking the term of lowest total degree. The expressions $\mathbf{e}$ and $\mathbf{a}^{\mathbf{e}}$ are sometimes referred to as "additive" versus "multiplicative" notation in the literature.

The *Hilbert series* of a finitely generated, graded S-module M is the expression

$$H(M; \mathbf{a}) = \sum_{\mathbf{e} \in \mathbb{Z}^d} \dim_K M_{\mathbf{e}} \mathbf{a}^{\mathbf{e}}$$

in the additive group $\prod_{\mathbf{e} \in \mathbb{Z}^d} \mathbb{Z}\mathbf{a}^{\mathbf{e}}$. Elements of this group are often called *Laurent series* in $\mathbf{a}$. The *K-polynomial* of M can be defined as

$$\mathcal{K}_S(M; \mathbf{a}) = \frac{H(M; \mathbf{a})}{H(S; \mathbf{a})}$$

where S is considered as a module over itself on the right hand side. This ratio of Laurent series is actually a Laurent polynomial. The K-polynomial can also be computed from a multigraded free resolution of M. In this paper we are most interested in modules $M = S/I$ where I is a homogeneous ideal in S; in this case, we sometimes write $\mathcal{K}_X(Y; \mathbf{a})$ for $\mathcal{K}_S(S/I; \mathbf{a})$, where $Y = \mathrm{Spec}(S/I) \subseteq X = \mathrm{Spec}(S)$. Also note that if we extend I to a larger polynomial ring $S[x]$, then the K-polynomial of S/I does not change.

The *multidegree* $\mathcal{C}(M; \mathbf{a})$ is obtained by substituting $1 - a_i$ for each variable a_i, substituting $(1 - a_i)^{-1} = \sum_{j \geq 0} a_i^j$ as necessary to get positive exponents, then taking the terms of lowest total degree. It generalizes the classical notion of degree to the multigraded setting, and when $M = S/I$, it can be geometrically interpreted as the class of $\mathrm{Spec}(S/I)$ in the equivariant Chow ring $A_T^*(\mathrm{Spec}(S))$.

Definition 2.25. The *K-theoretic quiver polynomial* $K\mathcal{Q}_{\mathbf{r}}(\mathbf{t}/\mathbf{s})$ (resp., *quiver polynomial* $\mathcal{Q}_{\mathbf{r}}(\mathbf{t} - \mathbf{s})$) is the K-polynomial (resp., multidegree) of the orbit closure $\Omega_{\mathbf{r}}$ with respect to its embedding in `rep` and multigrading described in §2.1. □

Our formulas for these polynomials will be in terms of the Grothendieck polynomials and Schubert polynomials of Lascoux and Schützenberger [LS82], which we briefly review now, following the definitions in [KM05]. Let $\mathbf{a} = (a_1, a_2, \dots)$ and $\mathbf{b} = (b_1, b_2, \dots)$ be alphabets and w_0 the longest element of the symmetric group S_m. We write $\ell(v)$ for the length of $v \in S_m$. The "top" double Grothendieck polynomial is defined as

$$\mathfrak{G}_{w_0}(\mathbf{a}; \mathbf{b}) = \prod_{i+j \leq m} \left(1 - \frac{a_i}{b_j}\right).$$

The *Demazure operator* $\overline{\partial_i}$ acts on polynomials in $\mathbf{a}$ by

$$\overline{\partial_i} f(a_i, a_{i+1}) = \frac{a_{i+1} f(a_i, a_{i+1}) - a_i f(a_{i+1}, a_i)}{a_{i+1} - a_i}$$

where f is written as a polynomial in only a_i, a_{i+1} with the other variables considered as coefficients. With this, we can inductively define the *double Grothendieck polynomial* of a permutation $ws_i \in S_m$ as $\mathfrak{G}_{ws_i}(\mathbf{a}; \mathbf{b}) = \overline{\partial_i} \mathfrak{G}_w(\mathbf{a}; \mathbf{b})$ whenever $\ell(ws_i) < \ell(w)$. This differs slightly from the definition in [FL94]; their Grothendieck polynomials $G_w(\mathbf{a}; \mathbf{b})$ are related to ours by the substitution $G_w(\mathbf{a}; \mathbf{b}) = \mathfrak{G}_w(\mathbf{a}^{-1}; \mathbf{b}^{-1})$, where an $\mathbf{a}^{-1} = (a_1^{-1}, a_2^{-1}, \dots)$ and similarly for $\mathbf{b}^{-1}$. The *double Schubert polynomial* $\mathfrak{S}_v(\mathbf{a}; \mathbf{b})$ of a permutation v is obtained from $\mathfrak{G}_v(\mathbf{a}; \mathbf{b})$ by the same process that a multidegree is obtained from a K-polynomial. We do not define Grothendieck polynomials associated to partial permutations, as is done in some literature, because we need two different ways of completing a partial permutation to an honest permutation (see §2.5). Instead, we give the explicit type of completion in the subscript of each occurrence to avoid ambiguity, at the expense of a slightly bulkier notation.

This gives a combinatorial construction of K-polynomials of matrix Schubert varieties. Recall that $\widetilde{\mathbf{a}}$ denotes the alphabet $\mathbf{a}$ written in reverse order. The following theorem was proven in the language of degeneracy loci in [Buc02, Theorem 2.1]; the formulation below is [KM05, Theorem A]. The multidegree version appeared in [Ful92].

Theorem 2.26. *Let $X = Mat(m, m)$, let $v \in S_m$, and let w_0 denote the longest element in S_m. The K-polynomials of matrix Schubert varieties, with the multigrading as described in §2.5, are the following Grothendieck polynomials:*

$$\mathcal{K}_X(X_v; \mathbf{a}, \mathbf{b}) = \mathfrak{G}_v(\mathbf{a}; \mathbf{b}) \qquad \textit{and} \quad \mathcal{K}_X(X^v; \mathbf{a}, \mathbf{b}) = \mathfrak{G}_{w_0 v w_0}(\widetilde{\mathbf{a}}; \widetilde{\mathbf{b}}). \tag{2.27}$$

The multidegrees of matrix Schubert varieties are the corresponding double Schubert polynomials.

Remark 2.28. In this paper, we often use K-polynomials and multidegrees of matrix Schubert varieties associated to arbitrary partial permutation matrices. The above theorem gives these as well: let v be $k \times l$ partial permutation matrix, let $c(v) \in S_m$ denote its completion, and let $oc(v) \in S_n$ denote its opposite completion (see §2.5). Then we have that

$$\mathcal{K}_{\mathrm{Mat}(k,l)}(\mathrm{Mat}(k,l)_v; \mathbf{a}, \mathbf{b}) = \mathcal{K}_{\mathrm{Mat}(m,m)}(\mathrm{Mat}(m,m)_{c(v)}; \mathbf{a}, \mathbf{b}) \qquad \text{and} \tag{2.29}$$

$$\mathcal{K}_{\mathrm{Mat}(k,l)}(\mathrm{Mat}(k,l)^v; \mathbf{a}, \mathbf{b}) = \mathcal{K}_{\mathrm{Mat}(n,n)}(\mathrm{Mat}(n,n)^{oc(v)}; \mathbf{a}, \mathbf{b}). \tag{2.30}$$

The analogous statements hold for multidegrees. These equations follow from the fact that the ideals I_v and $I_{c(v)}$ can be generated by the same polynomials, and similarly for I^v and $I^{oc(v)}$ (see §2.5). □



FIGURE 3. Two minimal lacing diagrams for the same orbit and their extended lacing diagrams.

2.8. **Lacing diagrams.** Lacing diagrams were introduced in [ADF85] to visualize type A quiver representations, and were interpreted as sequences of partial permutation matrices in [KMS06]. In this section we recall the essentials of lacing diagrams in arbitrary orientation. Our conventions differ from those used in [BR07] in order to make the connection with pipe dreams in §2.10 more natural. A *lacing diagram* for a type A quiver Q consists of columns of dots with arrows between certain pairs. The columns are indexed by the vertices of Q, and the number of dots in a column is the value of $\mathbf{d}$ at the corresponding vertex.

We use the following convention for vertically aligning the dots: if the columns are connected by an arrow pointing to the left, then the dots are aligned at the bottom; if the arrows connecting the columns points right, then the dots are aligned at the top. Then arrows can be placed between dots in the direction of the corresponding arrow of Q, with the restriction that each dot can be connected to at most one dot in each of its adjacent columns. Two examples of lacing diagrams continuing the running example are on the left side of Figure 3.

A lacing diagram can be interpreted as a sequence of partial permutation matrices $\mathbf{w} = (w_a)_{a \in Q_1}$. For each arrow a, the partial permutation matrix w_a has a 1 in row i, column j whenever there is an arrow from the ith dot from the top of the source column to the jth dot from the top of the target column. So, there is a 1 in location (i, j) in w_a if and only if $\vec{e}_i w_a = \vec{e}_j$ where $\vec{e}_i$ denotes the i^{th} standard basis (row) vector in the source vector space V_{ha}, and $\vec{e}_j$ denotes the j^{th} standard basis (row) vector in the target vector space V_{ta}. Since the sequence of partial permutation matrices $\mathbf{w}$ encodes exactly the same data as a lacing diagram, we will simply refer to $\mathbf{w}$ as a lacing diagram as well.

A lacing diagram can be completed to an *extended lacing diagram* using the completions and opposite completions of partial permutation matrices from §2.5 as follows: if w is a partial permutation matrix associated to a right pointing arrow, replace w with its completion $c(w)$, and if w is a partial permutation matrix associated to a left-pointing arrow, replace w by its opposite completion $oc(w)$. An extended lacing diagram can be visualized by adding “virtual” red dots and arrows to the original lacing diagram as in the right of Figure 3. The length $|\mathbf{w}|$ of a lacing diagram $\mathbf{w}$ is defined as the sum of the lengths of the permutations in the extended lacing diagram, or equivalently, the total number of crossings of laces in the extended lacing diagram.

Note that a lacing diagram $\mathbf{w}$ is naturally an element of `rep` by assigning the matrix w_a to each arrow $a \in Q_1$. The visualization as a lacing diagram allows one to easily see the indecomposable direct summands of this representation (the laces). Thus, two lacing diagrams $\mathbf{w}, \mathbf{w}'$ lie in the same GL-orbit if and only if they have the same number of laces between each pair of columns. A *minimal lacing diagram* is one whose length is minimal among those in its GL-orbit, and we denote by $W(\mathbf{r})$ the set of all minimal lacing diagrams in the orbit with rank array $\mathbf{r}$. For any $\mathbf{w} \in W(\mathbf{r})$, we have that the codimension of $\Omega_{\mathbf{r}}$ is $|\mathbf{w}|$ [BR07, Corollary 2].

It is known from [BR07, Prop. 1] that two lacing diagrams lie in the same orbit if and only if they are related by a series of transformations of the following form,

(2.31)



where both middle dots and at least one dot in each of the outer columns is not virtual (otherwise one of the permutations would not be a minimal length extension of the original partial permutation). There are also K-theoretic transformations of lacing diagrams

(2.32)



with the same condition on the dots, and in addition the two middle dots should be consecutive in their column. A *K-theoretic lacing diagram* for an orbit is one that can be obtained by these K-theoretic transformations from a minimal lacing diagram for the orbit. We let $KW(\mathbf{r})$ denote the set of K-theoretic lacing diagrams for the orbit with rank array $\mathbf{r}$.

The next theorem gives a characterization of the Zelevinsky permutation of an orbit directly from any lacing diagram in the orbit, analogous to [KMS06, Prop. 1.6].

Theorem 2.33. *Assume Q is a bipartite type A quiver and recall the indexing of blocks of a $d \times d$ matrix by vertices of Q from §2.4. Let $\mathbf{w}$ be any lacing diagram with quiver rank array $\mathbf{r}$. Then the number of 1s in $v(\mathbf{r})$ in the block with row indexed by vertex z_i and column indexed by vertex z_j is equal to:*

(Z1) the number of laces with left endpoint z_i and right endpoint z_j, whenever this makes sense;

(Z2) the number of arrows between the columns indexed by z_i and z_j whenever (z_i, z_j) is of the form (y_{k-1}, x_k) or (x_k, y_k) (i.e., z_i is exactly one vertex to the right of z_j);

(Z3) zero otherwise.

Proof. Let M be the matrix determined by $\mathbf{w}$ from (Z1)–(Z3), with the condition that the 1s of M are arranged from northwest to southeast in each block row and block column. To see that M is $v(\mathbf{r})$, it is enough to show that they have the same number of 1s in each block, since the arrangement of 1s in each block is then uniquely determined. First we reduce to checking this in the case that $\mathbf{w}$ contains a single lace. The number of 1s in each block of M is additive with respect to direct sum of representations (adding laces) by the definition. The number of 1s in each block of $v(\mathbf{r})$ is additive with respect to direct sum of representations because it is by definition a linear function of northwest block ranks $\mathbf{b}(\mathbf{r})_{i,j}$, which are in turn linear functions of the entries of the quiver rank array $\mathbf{r}$ [KR15, Lemma A.3]. The latter are additive with respect to direct sum of representations because they are, up to constants determined by $\mathbf{d}$, the dimensions of certain Hom spaces in the category of representations of Q (see the proof of [KR15, Prop. 3.1]).

Now we can assume $\mathbf{w}$ consists of a single lace. Noting that all blocks are of size one in this case, we will show that M and $\zeta(\mathbf{w})$ have the same number of 1s in each block by showing that they have the same northwest block ranks, which is equivalent to being in the same $B_- \times B_+$-orbit. This involves a slightly different computation for each of the 4 cases of the endpoints being x versus y type. An example of both M and $\zeta(\mathbf{w})$ is shown below for Q as in (2.4), with $\mathbf{w}$ the lace from y_3 to y_0.

$$M = \left[\begin{array}{ccc|cccc} 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{array}\right] \qquad \zeta(\mathbf{w}) = \left[\begin{array}{ccc|cccc} 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ \hline 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{array}\right] \tag{2.34}$$

Rather than introducing sufficiently complicated notation to formalize the computation, we carry out the computation in steps for this example. Since each step is just multiplication by a unipotent matrix with a region of alternating ±1s, it unambiguously generalizes to arbitrary size.

First, clear extra 1s from the northwest quadrant of $\zeta(\mathbf{w})$ by downwards row operations given by the following element of B_-:

$$\left[\begin{array}{ccccccc} 1 & & & & & & \\ -1 & 1 & & & & & \\ 1 & -1 & 1 & & & & \\ -1 & 1 & -1 & 1 & & & \\ \hline 0 & 0 & 0 & 0 & 1 & & \\ 0 & 0 & 0 & 0 & 0 & 1 & \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{array}\right] \cdot \zeta(\mathbf{w}) = \left[\begin{array}{ccc|cccc} 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 & -1 & 1 \\ \hline 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{array}\right] := M_1. \tag{2.35}$$

Now clear the top three rows of the northeast quadrant using column operations given by the following element of B_+:

$$M_1 \cdot \left[\begin{array}{ccc|cccc} 1 & 0 & 0 & -1 & 1 & -1 & 0 \\ & 1 & 0 & 1 & -1 & 0 & 0 \\ & & 1 & -1 & 0 & 0 & 0 \\ & & & 1 & 0 & 0 & 0 \\ & & & & 1 & 0 & 0 \\ & & & & & 1 & 0 \\ & & & & & & 1 \end{array}\right] = \left[\begin{array}{ccc|cccc} 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & -1 & 1 \\ \hline 1 & 0 & 0 & -1 & 1 & -1 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 & 0 \end{array}\right] := M_2. \tag{2.36}$$

The top three rows now clear the identity matrix in the southwest quadrant in an obvious way, so we omit the corresponding element of B_-, resulting in

$$\left[\begin{array}{ccc|cccc} 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & -1 & 1 \\ \hline 0 & 0 & 0 & -1 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \end{array}\right] := M_3. \tag{2.37}$$

Row operations given by the following element of B_- simplify the southeast quadrant further:

$$\left[\begin{array}{ccccccc} 1 &&&&&& \\ 0 & 1 &&&&& \\ 0 & 0 & 1 &&&& \\ 0 & 0 & 0 & 1 &&& \\ \hline 0 & 0 & 0 & 1 & 1 && \\ 0 & 0 & 0 & -1 & 0 & 1 & \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \end{array}\right] \cdot M_3 = \left[\begin{array}{ccc|cccc} 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & -1 & 1 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & -1 & 1 & -1 \end{array}\right] := M_4. \tag{2.38}$$

Now it is clear that M is the unique permutation matrix in this $B_- \times B_+$-orbit, by switching signs of the bottom 4 rows then using the leftmost 1 in each column to clear to the right, starting with column 4 and proceeding to the right. This completes the computation in the case of two y-type endpoints. The other three cases of the endpoints of $\mathbf{w}$ being x- or y-type are similar. ☐

Example 2.39. The lacing diagrams in Figure 3 each have 4 laces, and the number of arrows between adjacent columns of dots can also be readily seen, so one can apply the lemma above to produce the Zelevinsky permutation in (2.19) without calculating any northwest block ranks. ☐

Besides the GL-orbit of a lacing diagram $\mathbf{w} \in \texttt{rep}$, we will also be interested in a certain product of matrix Schubert varieties corresponding to the constituents of $\mathbf{w}$. For any arbitrary type A quiver Q, we define a closed subvariety of `rep`

$$\mathcal{O}(\mathbf{w}) = \prod_{\xrightarrow{a}} \mathrm{Mat}(\mathbf{d}(ta), \mathbf{d}(ha))_{w_a} \times \prod_{\xleftarrow{a}} \mathrm{Mat}(\mathbf{d}(ta), \mathbf{d}(ha))^{w_a}. \tag{2.40}$$

where the first product is taken over right pointing arrows of Q and the second over left pointing arrows. Notice that the codimension of $\mathcal{O}(\mathbf{w})$ in `rep` is $|\mathbf{w}|$.

2.9. Pipe dreams.

Our formulas use the language of pipe dreams, an introduction to which can be found in [MS05, §16.1]. These are the same as the RC-graphs originally introduced by Bergeron and Billey [BB93], which are based on the pseudo-line arrangements introduced by Fomin and Kirillov to the study of Grothendieck and Schubert polynomials [FK96]. The papers [BR04, WY12] also give helpful treatments of the topic.

Consider a $k \times l$ grid of squares. We use standard matrix terminology to refer to positions on the grid: horizontal strips of the squares are rows, vertical strips of squares are columns, and the square in row i from the top and column j from the left is labeled (i, j). A *pipe dream* on this grid is a subset of $\{1, \dots, k\} \times \{1, \dots, l\}$. A pipe dream P is visualized by tiling the grid using two kinds of tiles. For each (i, j) in the grid, place the corresponding tile according to the rule below:

$+$ (cross) if $(i, j) \in P$, (elbow) if $(i, j) \notin P$.

Let $|P|$ denote the cardinality of the subset P, which visually corresponds to the number of $+$ tiles. For a pipe dream P, denote by P_{rot} the pipe dream obtained by rotating the grid 180°. Two examples of pipe dreams on a 7×7 grid are in Figure 4; we have $|P_1| = 27$ and $|P_2| = 28$. The row and column labels come from identification of this grid with the northwest quadrant of $Y_\circ^{v_0}$ in the running example, with rows and columns labeled as in Figure 2.

A pipe dream P determines a word in the Coxeter generators $s_1, s_2, \dots$ of the symmetric group on $\mathbb{Z}_{>0}$, as follows. Starting from the top right of the grid, reading first along each row from right to left, then proceeding down the rows, one writes the letter s_{i+j-1} whenever



FIGURE 4. Pipe dreams related to the running example

one encounters a + in position (i, j). The word is written left to right. Evaluating this word using the relations

$$s_i^2 = s_i, \qquad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}, \qquad s_i s_j = s_j s_i, \text{ for } |i-j| \geq 2 \tag{2.41}$$

yields a permutation, known as the *Demazure product* of P, denoted $\delta(P)$. This is the same, up to sign, as taking the product in the degenerate Hecke/Iwahori algebra [FK94, FK96]. For a given permutation v, we say that P is a *pipe dream for* v if $\delta(P) = v$. If P is a pipe dream such that $|P| = \ell(\delta(P))$, then we say that P is *reduced*. Both pipe dreams in Figure 4 have Demazure product equal to the Zelevinsky permutation in (2.19). The one of the left is reduced, the other is not because $|P_2| = 28$ while $\delta(P) = 27$.

The relation between P and $\delta(P)$ has the visual interpretation that $\delta(P)$ can be read off by "following" the pipes. First consider the case P is reduced, which occurs if and only if no pair of pipes crosses more than once. We can expand the grid to a large enough $n \times n$ square, and consider $\delta(P)$ as a permutation of $\{1, 2, \ldots, n\}$. Then, for each $i \in \{1, 2, \ldots, n\}$, there is a pipe in P connecting row i on the left to column $\delta(P)(i)$ at the top. Now for an arbitrary P, we do the same except that for each pair of pipes we only allow them to cross once and ignore any additional crossings (i.e., the visualization above will hold for any reduced sub-pipe dream $P' \subset P$ such that $\delta(P') = \delta(P)$). In the non-reduced pipe dream P_2 of Figure 4, we see two pipes towards the lower left crossing twice.

Finally, we need a relative version of pipe dreams which uses another permutation to constrain where + tiles are allowed to be placed. Recall from §2.5 the diagram $\mathcal{D}_w$ of a partial permutation w. For two permutations v, w, denote by $\operatorname{Pipes}(w, v)$ the set of pipe dreams P for v such that $P \subseteq \mathcal{D}_w$, and similarly define $\operatorname{RedPipes}(w, v)$ to consist of the reduced pipe dreams in $\operatorname{Pipes}(w, v)$. The pipe dreams in Figure 4 are pipe dreams on the diagram of v_0 from (2.11).

Fix a grid which will be tiled to produce a pipe dream. Let $\mathbf{a}$ and $\mathbf{b}$ be alphabets indexing the rows and columns of the grid, respectively. For a pipe dream P, define

$$(\mathbf{1} - \mathbf{a}/\mathbf{b})^P = \prod_{(i,j)\in P} (1 - a_i/b_j)$$

Similarly, define

$$(\mathbf{a} - \mathbf{b})^P = \prod_{(i,j)\in P} (a_i - b_j).$$

We use the following formulas for K-polynomials of Kazhdan-Lusztig varieties, as presented in [WY12, Theorem 4.5]. These formulas previously appeared in other forms in [AJS94, Bil99] and [Gra02, Wil06].

Theorem 2.42. *Recall from §2.6 the multigrading of $Y_\circ^{v_0}$ by the free abelian group on alphabets $\mathbf{s}, \mathbf{t}$. The K-polynomial of the Kazhdan-Lusztig variety Y_v inside $Y_\circ^{v_0}$ is given by each of the following two formulas:*

$$\mathcal{K}_{Y_\circ^{v_0}}(Y_v; \mathbf{s}, \mathbf{t}) = \sum_{P \in \mathrm{Pipes}(v_0, v)} (-1)^{|P| - l(v)} (\mathbf{1} - \mathbf{t}/\mathbf{s})^P \tag{2.43}$$

$$\mathcal{K}_{Y_\circ^{v_0}}(Y_v; \mathbf{s}, \mathbf{t}) = \mathfrak{G}_v(\mathbf{t}, \mathbf{s}; \mathbf{s}, \mathbf{t}), \tag{2.44}$$

and its multidegree is given by each of:

$$\mathcal{C}_{Y_\circ^{v_0}}(Y_v; \mathbf{s}, \mathbf{t}) = \sum_{P \in \mathrm{RedPipes}(v_0, v)} (\mathbf{t} - \mathbf{s})^P \tag{2.45}$$

$$\mathcal{C}_{Y_\circ^{v_0}}(Y_v; \mathbf{s}, \mathbf{t}) = \mathfrak{S}_v(\mathbf{t}, \mathbf{s}; \mathbf{s}, \mathbf{t}). \tag{2.46}$$

2.10. Relating pipes and laces.

We need a connection between pipe dreams and lacing diagrams for the degeneration construction in §4. This will be for bipartite quivers, so we restrict our consideration to these. We first show that all pipe dreams in $\mathrm{Pipes}(v_0, v(\mathbf{r}))$ contain the following particular pipe dream as a subdiagram: define P_* to be the pipe dream on the Rothe diagram of v_0, which has a $+$ tile in location (i,j) if and only if (i,j) lies outside the snake region (Definition 2.13 and Figure 1).

Lemma 2.47. *Every element of* $\mathrm{Pipes}(v_0, v(\mathbf{r}))$ *contains P_* as a subdiagram. Furthermore* $\mathrm{Pipes}(v_0, v_*) = \{P_*\}$*, where v_* denotes the Zelevinsky permutation associated to the entire representation space* `rep`.

Proof. Let $I_{v(\mathbf{r})} \subseteq K[Y_\circ^{v_0}]$ denote the Kazhdan-Lusztig ideal that defines the Zelevinsky image $\zeta(\Omega_\mathbf{r}) = Y_{v(\mathbf{r})}$, and let $\prec$ be a lexicographical antidiagonal monomial order on the variables in $K[Y_\circ^{v_0}]$ as in [WY12]. For each $(i,j) \in P_*$, the variable $z_{ij} \in I_{v(\mathbf{r})}$ since the corresponding matrix position is always zero for $Y_{v(\mathbf{r})}$, and thus also $z_{ij} \in \mathrm{in}_\prec I_{v(\mathbf{r})}$ since z_{ij} is just a monomial. Now, by [WY12, Theorem 3.2], we have

$$\mathrm{in}_\prec I_{v(\mathbf{r})} = \bigcap_{P \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))} \langle z_{ij} \mid (i,j) \in P \rangle, \tag{2.48}$$

So $(i,j) \in P$ for every $P \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))$, and consequently, every such P must contain P_*. Since every (possibly non-reduced) pipe dream must contain a reduced pipe dream as a subdiagram, we obtain the first statement of the lemma.

For the second statement, observe that I_{v_*} is the monomial ideal generated by the variables outside of the snake region. Consequently, P_* is the unique element of $\mathrm{RedPipes}(v_0, v_*)$. But every element of $\mathrm{Pipes}(v_0, v(\mathbf{r}))$ is contained in the union of elements of $\mathrm{RedPipes}(v_0, v(\mathbf{r}))$ by [Mil05, Lemma 2], so we see that $\mathrm{Pipes}(v_0, v(\mathbf{r})) = \{P_*\}$. ☐

To define the lacing diagram $\mathbf{w}(P)$ determined by a pipe dream P, we first consider the case of $P \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))$. In this case, we can simply follow the pipes through each block of the snake region: to be precise, consider the block corresponding to an arrow α_i of Q. For each pipe passing through the right and bottom boundaries of this block, there is a 1 in the row and column of the partial permutation matrix w_{α_i} corresponding to the row and column where the pipe meets these boundaries, and the rest of the matrix is filled with 0s. For the block corresponding to β_i, a pipe passing through the top and left boundaries contributes a 1 in the row and column of w_{β_i} corresponding to the row and column where the pipe meets these boundaries, and the rest is filled with 0s. Now for arbitrary $P \in \mathrm{Pipes}(v_0, v(\mathbf{r}))$, let P^{α_i} be the restriction of P to the block of the snake region indexed by α_i, and similarly for P^{β_i}. Take a reduced pipe dream $P' \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))$

such that $P' \subseteq P$, and also both $\delta(P^{\beta_i}) = \delta({P'}^{\beta_i})$ and $\delta(P^{\alpha_i}_{\mathrm{rot}}) = \delta({P'}^{\alpha_i}_{\mathrm{rot}})$. Then we define $\mathbf{w}(P) = \mathbf{w}(P')$, which does not depend on P'.

Example 2.49. For the reduced pipe dream P_1 from Figure 4, we follow the pipes in each block and find that $\mathbf{w}(P_1)$ is the top lacing diagram of Figure 3. Notice that some pipe crossings are only visible in the extended diagram. For the non-reduced P_2, we have that $P_1 \subset P_2$ and they have the same Demazure product within each block (after rotation for type α blocks), so $\mathbf{w}(P_2) = \mathbf{w}(P_1)$. □

Proposition 2.50 below gives a connection between certain reduced pipes dreams and minimal lacing diagrams. This is the bipartite generalization of both [KMS06, Theorem 5.10] and [KMS06, Corollary 6.18]. We first make some preliminary observations about the relation between P and $\mathbf{w}(P)$. The reader may find the row and column labels in Figure 1 and examples of pipe dreams in Figure 4 helpful for visualization. Identify the diagram of v_0 with the northwest quadrant of $Y_\circ^{v_0}$, keeping in mind the labeling of the block rows and columns of $Y_\circ^{v_0}$ by vertices of Q from §2.4. For a fixed $P \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))$, each individual pipe of P has an associated row and column vertex determined by the labels of the row and column blocks of the endpoints of the pipe (the area outside the diagram of v_0, i.e. the outside northwest quadrant of $Y_\circ^{v_0}$, is filled with elbow tiles). Then, we can see that:

(i) every pipe passes through the snake region;
(ii) if two pipes cross outside the snake region, then they cannot cross again;
(iii) the row and column vertices of a pipe can be read off by the walls of the snake which the pipe passes through.

Making (iii) more precise, we have the following cases.

y-x: A pipe has a y-type row vertex and an x-type column vertex if and only if it meets the snake boundary at a left wall and an upper wall.

y-y: A pipe has both row and column vertices y-type if and only if it meets the snake boundary at a left wall and a right wall.

x-y: A pipe has an x-type row vertex and a y-type column vertex if and only if it meets the snake boundary at a bottom wall and a right wall.

x-x: A pipe has both row and column vertices x-type if and only if it meets the snake boundary at a bottom wall and an upper wall.

Now we come to one of the key combinatorial ingredients of our K-theoretic component formula.

Proposition 2.50. *The following hold:*

(a) for any $P \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))$, we have that $\mathbf{w}(P) \in W(\mathbf{r})$;
(b) for any $\mathbf{w}_0 \in W(\mathbf{r})$, there exists a $P \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))$ such that $\mathbf{w}(P) = \mathbf{w}_0$.

Proof. (a) Let v' be the Zelevinsky permutation associated to the orbit closure of $\mathbf{w}(P)$. First we show that $v' = v(\mathbf{r})$ using Theorem 2.33. Since we know that v' is the Zelevinsky permutation of some orbit, we only need to compare the blocks in condition (Z1) because blocks in (Z2) and (Z3) contain redundant information. Let (z_i, z_j) be a pair of vertices with z_i weakly to the left of z_j. By the definition of $\mathbf{w}(P)$ and remarks in the paragraph preceding this proposition, the number of laces in $\mathbf{w}(P)$ from z_i to z_j is the number of pipes with row labeled by z_i at the left to the block column labeled by z_j at top. This is the same as for the corresponding block of $v(\mathbf{r})$ since P is reduced and $\delta(P) = v(\mathbf{r})$. So $\mathbf{w}(P)$ is a lacing diagram in the orbit indexed by $\mathbf{r}$.

Now the fact that $\mathbf{w}(P)$ is a minimal lacing diagram is guaranteed by the fact that

$$|\mathbf{w}(P)| = \#(+\text{ tiles in snake region of } P) = \operatorname{codim} \Omega_{\mathbf{r}} \text{ in } \mathtt{rep}, \tag{2.51}$$

where the first equality holds by definition of $\mathbf{w}(P)$ and the second holds by Lemma 2.47 together with the fact that P is a reduced pipe dream (so that the total number of +s in P is the codimension of $\zeta(\Omega_{\mathbf{r}})$ in $Y_\circ^{v_0}$).

(b) By simply choosing any placement of + tiles within the snake to make the pipes cross in the same configuration as $\mathbf{w}_0$, we see that there exist pipe dreams P on the diagram of v_0 with $\mathbf{w}(P) = \mathbf{w}_0$ such that $P \supseteq P_*$ and the restriction of P to any block is reduced. Fix such a P; we will show that any such P is reduced and $\delta(P) = v(\mathbf{r})$. That P is reduced is straightforward: if two pipes cross outside the snake region, they cannot cross again by observation (ii) above. So any double crossings must come from two pipes that cross twice within the snake region. But since $\mathbf{w}_0$ is minimal, no two laces cross twice, so no two pipes of P can cross twice in the snake region. So P is reduced.

To show that $\delta(P) = v(\mathbf{r})$, we again use the characterization of $v(\mathbf{r})$ in Theorem 2.33, and again just have to compare the blocks in condition (Z1), say a block with row index z_i and column index z_j where z_i is weakly to the left of z_j in Q. Since we have shown P is reduced, $\delta(P)$ can be read off by following the pipes. The number of 1s in this block of $\delta(P)$ is the number of pipes running between block row z_i and block column z_j, which is equivalent to knowing the number of pipes running between certain boundaries of the snake region by the observations preceding the proposition (there are four cases depending on which of z_i, z_j is x-type and y-type). But having chosen $\mathbf{w}(P) = \mathbf{w}_0$, we have made it so that in each case this is exactly the number of laces with left endpoint z_i and right endpoint z_j, which is the same number of 1s in the corresponding block of $v(\mathbf{r})$ by Theorem 2.33, since $\mathbf{w}_0$ is assumed to be a lacing diagram with rank array $\mathbf{r}$. □

2.11. **Gröbner degenerations and flat families.** We use *Gröbner degenerations* in our proof of the component formulas. Here we briefly summarize the relevant background, referring the reader to the textbook [Eis95, §15.8] for a similar treatment in more detail.

An *integral weight function* $\lambda\colon \mathbb{Z}^n \to \mathbb{Z}$ is a homomorphism of groups used to assign an integer value $\lambda(\mathbf{a})$ (called a *weight*) to each monomial $\mathbf{x}^{\mathbf{a}} = x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n} \in K[x_1, \ldots, x_n]$. Sometimes we will write $\lambda(\mathbf{x}^{\mathbf{a}})$ instead of $\lambda(\mathbf{a})$ to denote the weight of the monomial $\mathbf{x}^{\mathbf{a}}$. The *initial form* $\mathrm{in}_\lambda(f)$ of a polynomial $f \in K[x_1, \ldots, x_n]$ with respect to the integral weight function λ is the sum of the terms of f of highest weight. For example, if x_1 and x_2 in $K[x_1, x_2]$ have weights 1 and 3 respectively, then the initial form of $f := x_1^3 - 5x_2 + x_1^2 + 2$ is $x_1^3 - 5x_2$, since x_1^3 and $-5x_2$ are the terms of highest weight in f. Define the *initial ideal* $\mathrm{in}_\lambda(I)$ of an ideal $I \subseteq K[x_1, \ldots, x_n]$ to be

$$\mathrm{in}_\lambda(I) := \langle \mathrm{in}_\lambda(f) \mid f \in I \rangle$$

A *Gröbner basis* $G = \{g_1, \ldots, g_r\}$ with respect to a choice of integral weight function λ is a generating set of I such that $\{\mathrm{in}_\lambda(g_1), \ldots, \mathrm{in}_\lambda(g_r)\}$ is a generating set of $\mathrm{in}_\lambda(I)$.

Given an integral weight function λ, one can *homogenize* an ideal $I \subseteq K[x_1, \ldots, x_n]$ to get an ideal $\widetilde{I} \in K[x_1, \ldots, x_n, t]$ defined as follows: let $f = \sum_k a_k m_k$, with $a_k \in K^\times$, m_k a monomial. Let $b(f)$ denote the max integer in the set $\{\lambda(m_k) \mid m_k \text{ is a monomial in } f\}$. Define

$$\widetilde{I} := \langle t^{b(f)} f(t^{-\lambda(x_1)} x_1, \ldots, t^{-\lambda(x_n)} x_n) \mid f \in I \rangle.$$

This is a slight abuse of notation since $\widetilde{I}$ depends on the choice of integral weight function. However, this is harmless as the integral weight function will always be fixed in advance. We will use the notation $\widetilde{I}(a)$ to denote the ideal of $K[x_1, \ldots, x_n]$ obtained by replacing the variable t with $a \in K$.

Theorem 2.52. *Let $I \subseteq K[x_1, \ldots, x_n]$ be a homogenous ideal with respect to the standard grading. Let $\lambda\colon \mathbb{Z}^n \to \mathbb{Z}$ be an integral weight function. Then, the following hold:*

(1) $K[x_1,\dots,x_n,t]/\widetilde{I}$ *is a flat* $K[t]$*-module;*
(2) the fiber over $t \neq 0$ *is isomorphic to* $K[x_1,\dots,x_n]/I$*;*
(3) the fiber over $t = 0$ *is* $K[x_1,\dots,x_n]/in_\lambda(I)$*.*

Remark 2.53. This theorem is a variation of [Eis95, Theorem 15.17], which does not assume that I is homogeneous, but whose proof assumes that the weight order induced by λ can be refined to a monomial order. To see that the theorem as stated above holds, let N be any integer so that $\lambda(x_i) + N > 0$ for all $1 \leq i \leq n$, and define a second weight function $\mu : \mathbb{Z}^n \to \mathbb{Z}$ with $\mu(x_i) = \lambda(x_i) + N$. Then, for any homogeneous polynomial f, $\mathrm{in}_\lambda(f) = \mathrm{in}_\mu(f)$, and so for a homogeneous ideal I, $\mathrm{in}_\lambda(I) = \mathrm{in}_\mu(I)$. One may similarly check that the λ-homogenization and μ-homogenization of a homogeneous (with respect to the standard grading of $K[x_1,\dots,x_n]$) ideal I are identical. Now since $\mu(x_i) > 0$ for all i, the weight order induced by μ can be refined to a monomial order by breaking ties $\mu(\mathbf{x}^\mathbf{a}) = \mu(\mathbf{x}^\mathbf{b})$ with a lexicographical order on the variables. So, by [Eis95, Theorem 15.17] the statement of the above theorem holds for the integral weight function μ and thus also for λ. □

An integral weight function is naturally associated to a coordinate $K^\times$-action on $\mathbb{A}^n$: if $t \in K^\times$ and $(a_1,\dots,a_n) \in \mathbb{A}^n$, then the action

$$t \cdot (a_1,\dots,a_n) = (t^{\lambda_1}a_1,\dots,t^{\lambda_n}a_n) \tag{2.54}$$

induces the integral weight function $\lambda\colon \mathbb{Z}^n \to \mathbb{Z}$ with $\lambda(x_i) = \lambda_i$. Consequently, one may define the *homogenization of an ideal* $I \subseteq K[x_1,\dots,x_n]$ *with respect to the weights from a* $K^\times$*-action* as the homogenization of I with respect to the associated integral weight function.

Let $X \subseteq \mathbb{A}^n$ denote the subscheme defined by $I \subseteq K[x_1,\dots,x_n]$, and let $\widetilde{X} \subseteq \mathbb{A}^n \times \mathbb{A}^1$ denote the subscheme defined by the homogenization of I with respect a coordinate $K^\times$-action (or, equivalently, its associated integral weight function). Denote this homogenization by $\widetilde{I} \subseteq K[x_1,\dots,x_n,t]$. By Theorem 2.52, $\widetilde{X} \subseteq \mathbb{A}^n \times \mathbb{A}^1$ is a flat family with fibers $\widetilde{X}(a) := \mathrm{Spec}(K[x_1,\dots,x_n,t]/\widetilde{I} \otimes_{K[t]} K[t]/\langle t-a\rangle)$, $a \in K$. If $a \neq 0$, the fiber $\widetilde{X}(a)$ is $a \cdot X$, for the given $K^\times$-action. The limit scheme $\lim_{t\to 0}(t \cdot X)$ is the fiber $\widetilde{X}(0)$, which has defining ideal $\widetilde{I}(0) = \mathrm{in}_\lambda I$. The scheme X is said to *Gröbner degenerate* to $\widetilde{X}(0)$, which we sometimes denote by $\mathrm{in}_\lambda X$.

Remark 2.55. We will also need Gröbner bases and degenerations for *monomial orders* (see [Eis95, §15] for background on this material). To fix notation, let $<$ be a monomial order on the polynomial ring $K[x_1,\dots,x_n]$. If $f \in K[x_1,\dots,x_n]$, then we use $\mathrm{in}_< f$ to denote the *initial term* of f with respect to the monomial order $<$. If I is an ideal in $K[x_1,\dots,x_n]$, then we use the notation $\mathrm{in}_< I$ to denote the *initial ideal*

$$\mathrm{in}_< I := \langle \mathrm{in}_< f \mid f \in I\rangle.$$

If $X = \mathrm{Spec}\, K[x_1,\dots,x_n]/I$, then we use $\mathrm{in}_< X$ to denote the *initial scheme*

$$\mathrm{in}_< X := \mathrm{Spec}\, K[x_1,\dots,x_n]/\mathrm{in}_< I,$$

and we say that X *Gröbner degenerates* to $\mathrm{in}_< X$. □

3. Bipartite ratio and pipe formulas

In this section and the next, the fixed bipartite quiver Q of type A and dimension vector $\mathbf{d}$ are omitted from the notation.

3.1. **Bipartite ratio formulas.** We include a proof of the ratio formula for completeness, though it follows easily from our construction of the bipartite Zelevinsky map using the same argument as the equioriented case [KMS06, Theorem 2.7].

Theorem 3.1 (Bipartite ratio formula). *For any rank array* $\mathbf{r}$*, we have*

$$KQ_{\mathbf{r}}(\mathbf{t}/\mathbf{s}) = \frac{\mathfrak{G}_{v(\mathbf{r})}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})}{\mathfrak{G}_{v_*}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})} \qquad \text{and} \qquad Q_{\mathbf{r}}(\mathbf{t}-\mathbf{s}) = \frac{\mathfrak{S}_{v(\mathbf{r})}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})}{\mathfrak{S}_{v_*}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})}. \tag{3.2}$$

Proof. Let $H(\Omega_{\mathbf{r}};\mathbf{s},\mathbf{t})$ denote the multigraded Hilbert series of $\Omega_{\mathbf{r}}$ for the multigrading described in §2.6. We have

$$H(\Omega_{\mathbf{r}};\mathbf{s},\mathbf{t}) = \mathcal{K}_{\mathtt{rep}}(\Omega_{\mathbf{r}};\mathbf{s},\mathbf{t})H(\mathtt{rep};\mathbf{s},\mathbf{t}), \tag{3.3}$$

and

$$H(\Omega_{\mathbf{r}};\mathbf{s},\mathbf{t}) = H(\zeta(\Omega_{\mathbf{r}});\mathbf{s},\mathbf{t}) = \mathcal{K}_{Y_\circ^{v_0}}(\zeta(\Omega_{\mathbf{r}});\mathbf{s},\mathbf{t})H(Y_\circ^{v_0};\mathbf{s},\mathbf{t}), \tag{3.4}$$

and

$$H(\mathtt{rep};\mathbf{s},\mathbf{t}) = \mathcal{K}_{Y_\circ^{v_0}}(\zeta(\mathtt{rep});\mathbf{s},\mathbf{t})H(Y_\circ^{v_0};\mathbf{s},\mathbf{t}). \tag{3.5}$$

Rearranging these equations yields

$$\mathcal{K}_{\mathtt{rep}}(\Omega_{\mathbf{r}};\mathbf{s},\mathbf{t}) = \frac{\mathcal{K}_{Y_\circ^{v_0}}(\zeta(\Omega_{\mathbf{r}});\mathbf{s},\mathbf{t})}{\mathcal{K}_{Y_\circ^{v_0}}(\zeta(\mathtt{rep});\mathbf{s},\mathbf{t})}. \tag{3.6}$$

Finally, because $\zeta(\Omega_{\mathbf{r}}) = Y_{v(\mathbf{r})}$, and $\zeta(\mathtt{rep}) = Y_{v_*}$, we can apply (2.44) to obtain the K-theoretic formula. The multidegree formula follows from the usual relationship between K-polynomials and multidegrees on both sides. □

3.2. **Bipartite pipe formulas.** Next, we provide formulas in terms of pipe dreams. To begin, recall that $P_* \in \text{RedPipes}(v_0, v_*)$ is the pipe dream that has a $+$ tile in location (i,j) if and only (i,j) lies outside of the snake region. We have:

Theorem 3.7 (Bipartite pipe formula). *For any rank array* $\mathbf{r}$*, we have*

$$KQ_{\mathbf{r}}(\mathbf{t}/\mathbf{s}) = \sum_{P\in\text{Pipes}(v_0,v(\mathbf{r}))} (-1)^{|P|-l(v(\mathbf{r}))}(\mathbf{1}-\mathbf{t}/\mathbf{s})^{P\setminus P_*} \tag{3.8}$$

and

$$Q_{\mathbf{r}}(\mathbf{t}-\mathbf{s}) = \sum_{P\in\text{RedPipes}(v_0,v(\mathbf{r}))} (\mathbf{t}-\mathbf{s})^{P\setminus P_*}. \tag{3.9}$$

Proof. First consider the K-theoretic version. We can express the numerator and denominator of (3.6) from the proof of the ratio formula as sums over pipe dreams using (2.43). Applying Lemma 2.47, we find that the denominator has only one term, and get

$$KQ_{\mathbf{r}}(\mathbf{t}/\mathbf{s}) = \frac{\sum_{P\in\text{Pipes}(v_0,v(\mathbf{r}))}(-1)^{|P|-l(v(\mathbf{r}))}(\mathbf{1}-\mathbf{t}/\mathbf{s})^{P}}{(\mathbf{1}-\mathbf{t}/\mathbf{s})^{P_*}}. \tag{3.10}$$

But also by Lemma 2.47 we know that the denominator divides every summand in the numerator, yielding (3.8). The multidegree formula (3.9) follows by the usual process to obtaining multidegrees from K-polynomials, noting that the degree of a summand is determined by the number of +s in the corresponding pipe dream, so the lowest degree terms are exactly indexed by reduced pipe dreams. □

4. Degeneration and bipartite component formula

The goal of this section is to prove the K-theoretic component formula (Theorem 4.37). This is accomplished by degenerating a quiver locus to a reduced union of products of matrix Schubert varieties (Theorem 4.28), which leaves the K-polynomial invariant, and the K-polynomial of the latter can be calculated more easily. As usual, an arbitrary dimension vector $\mathbf{d}$ is fixed and thus omitted from the notation when possible.

4.1. Degeneration of quiver loci.

We start with the basic idea outlined at [KMS06, p. 238]. We give a flat family of group schemes $\widetilde{GL}$ whose general fiber is isomorphic to GL and a flat family of schemes $\widetilde{\mathtt{rep}}$ whose general fiber is isomorphic to $\mathtt{rep}$. We show that the first family acts fiberwise on the second, so that each quiver locus $\Omega_{\mathbf{r}}$ in a general fiber degenerates in the special fiber of $\widetilde{\mathtt{rep}}$ to a (possibly non-reduced) union of orbit closures of the special fiber of $\widetilde{GL}$, which are products of matrix Schubert varieties. Finally, we show that the degenerations are actually reduced using the bipartite Zelevinsky map of [KR15] and ideas in the paper [WY12], with components of the degenerations indexed by minimal lacing diagrams.

Family of groups. At each vertex z, fix a basis $\{e^z_1, e^z_2, \dots, e^z_{\mathbf{d}(z)}\}$ of $E_z = K^{\mathbf{d}(z)}$. So we have in this notation that $GL = \prod_{z\in Q_0} GL(E_z)$, with $GL(E_z)$ acting on the right of E_z, inducing our standard action of GL on

$$\mathtt{rep} = \bigoplus_{a\in Q_1} \operatorname{Hom}(E_{ta},\, E_{ha}) = \bigoplus_{a\in Q_1} E^*_{ta} \otimes E_{ha}$$

as in Section 2.1. We introduced the "doubled" base-change group

$$GL^2 = GL \times GL = \{(g^\alpha_z,\, g^\beta_z) \mid z \in Q_0\},$$

inside which we will construct the family $\widetilde{GL}$ mentioned above. The family is parametrized by $\mathbb{A}^1 = \operatorname{Spec} K[t]$, utilizing a $K^\times$ action where we identify $K^\times = \operatorname{Spec} K[t, t^{-1}] \subset \mathbb{A}^1$.

For each $z \in Q_0$, consider the diagonal one-parameter subgroup $\rho_z \colon K^\times \to GL(E_z)$ defined by

$$e^z_i \cdot \rho_z(t) = t^i e^z_i, \qquad t \in K^\times. \tag{4.1}$$

We use this notation to define a right action of $K^\times$ on GL^2 at x-type vertices (omitting their subscripts) by

$$(g^\beta_x,\, g^\alpha_x) \cdot t = \Big(\rho_x(t^{-1}) g^\beta_x \rho_x(t),\ \rho_x(t) g^\alpha_x \rho_x(t^{-1})\Big). \tag{4.2}$$

for $t \in K^\times$, and at y-type vertices by

$$(g^\alpha_y,\, g^\beta_y) \cdot t = \Big(\rho_y(t^{-1}) g^\alpha_y \rho_y(t),\ \rho_y(t) g^\beta_y \rho_y(t^{-1})\Big). \tag{4.3}$$

We let $\widetilde{GL}^\circ$ be the inverse image of the diagonal $\Delta \subset GL^2$ under the action map of the above action $GL^2 \times K^\times \to GL^2$, and $\widetilde{GL}$ the closure of $\widetilde{GL}^\circ$ in $GL^2 \times \mathbb{A}^1$, where we identify $K^\times \subset \mathbb{A}^1$.

Consider the subgroup $B_+ \times B_- \subseteq GL^2$ defined by taking the subgroup of upper-triangular matrices in each factor $GL^\beta_{x_k}$ and $GL^\alpha_{y_k}$, and the subgroup of lower-triangular matrices in each factor $GL^\alpha_{x_k}$ and $GL^\beta_{y_k}$. So, like GL^2, each element of $B_+ \times B_-$ associates two matrices to each vertex. Then we denote by $B_+ \times_T B_-$ the subgroup of $B_+ \times B_-$ consisting of elements such that the pair at each vertex has the same diagonal. Now proving the following lemma is similar to [KMS06, Lemma 4.2].

Lemma 4.4. *The family $\widetilde{GL} \subseteq GL^2 \times \mathbb{A}^1$ is a flat family of closed group subschemes of GL^2, with general fiber over points of $K^\times$ isomorphic to GL, and special fiber over $t = 0$ equal to $B_+ \times_T B_-$.*

Proof. We first address flatness. For each factor $GL^\star(E_z)$ of GL^2 (where $\star$ stands for α or β), let $(g_{ij}^{z,\star})$ be coordinates giving the matrix entries in our fixed basis. The induced right action of $K^\times$ on coordinates is

$$\begin{aligned} g_{ij}^{x_k,\alpha} \cdot t = t^{i-j} g_{ij}^{x_k,\alpha}, \qquad & g_{ij}^{x_k,\beta} \cdot t = t^{-(i-j)} g_{ij}^{x_k,\beta}, \\ g_{ij}^{y_k,\alpha} \cdot t = t^{-(i-j)} g_{ij}^{y_k,\alpha}, \qquad & g_{ij}^{y_k,\beta} \cdot t = t^{i-j} g_{ij}^{y_k,\beta}. \end{aligned} \tag{4.5}$$

The diagonal $\Delta \subset GL^2$ is defined by the ideal $I_\Delta = \langle g_{ij}^{z,\beta} - g_{ij}^{z,\alpha} \rangle$, and its homogenization $\widetilde{I}_\Delta$ from §2.11 with respect to the $K^\times$-action in (4.5) is

$$\begin{aligned} \widetilde{I}_\Delta = & \sum_{k=1}^{n} \langle g_{ij}^{x_k,\beta} - g_{ij}^{x_k,\alpha} t^{2i-2j} \mid i \geq j \rangle + \langle g_{ij}^{x_k,\beta} t^{2j-2i} - g_{ij}^{x_k,\alpha} \mid i \leq j \rangle \\ & + \sum_{i=0}^{n} \langle g_{ij}^{y_k,\alpha} - g_{ij}^{y_k,\beta} t^{2i-2j} \mid i \geq j \rangle + \langle g_{ij}^{y_k,\alpha} t^{2j-2i} - g_{ij}^{y_k,\beta} \mid i \leq j \rangle. \end{aligned} \tag{4.6}$$

Thus, by Theorem 2.52 we know that $\widetilde{I}_\Delta$ defines a flat subscheme of $\mathrm{Mat}^2 \times \mathbb{A}^1$, where

$$\mathrm{Mat} := \prod_{z \in Q_0} \mathrm{Mat}(\mathbf{d}(z), \mathbf{d}(z)),$$

and $\mathrm{Mat}^2 = \mathrm{Mat} \times \mathrm{Mat}$. Since GL^2 is an open subscheme of Mat^2, the subscheme $\widetilde{GL}$ defined by $\widetilde{I}_\Delta$ in $GL^2 \times \mathbb{A}^1$ is also flat.

It remains to show that each fiber is a closed group subscheme of GL^2. Equations (4.2) and (4.3) show that $K^\times$ acts on GL^2 by group automorphisms, so the fiber over each $t \neq 0$ isomorphic to $\widetilde{GL}(1) = GL$. From (4.6), the fiber over $t = 0$ in $\widetilde{GL}$ is defined by the ideal

$$\begin{aligned} \widetilde{I}_\Delta(0) = & \sum_{k=1}^{n} \langle g_{ij}^{x_k,\beta} \mid i > j \rangle + \sum_{k=1}^{n} \langle g_{ij}^{x_k,\alpha} \mid i < j \rangle + \sum_{k=1}^{n} \langle g_{ii}^{x_k,\beta} - g_{ii}^{x_k,\alpha} \rangle \\ & + \sum_{k=1}^{n} \langle g_{ij}^{y_k,\alpha} \mid i > j \rangle + \sum_{k=1}^{n} \langle g_{ij}^{y_k,\beta} \mid i < j \rangle + \sum_{k=1}^{n} \langle g_{ii}^{y_k,\beta} - g_{ii}^{y_k,\alpha} \rangle \end{aligned} \tag{4.7}$$

which is clearly the ideal of the closed subgroup $B_+ \times_T B_-$. ☐

Family of quiver loci. We now define a right action of $K^\times$ on `rep` which is compatible with the $K^\times$-action on $\widetilde{GL}$ above. Retaining the definition of ρ_z from (4.1) and again omitting vertex and arrow subscripts for readability, we define

$$V_\beta \cdot t = \rho_y(t) V_\beta \, \rho_x(t) \qquad \text{and} \qquad V_\alpha \cdot t = \rho_y(t^{-1}) V_\alpha \, \rho_x(t^{-1}). \tag{4.8}$$

Define $\widetilde{\Omega}_{\mathbf{r}}^\circ$ as the inverse image of $\Omega_{\mathbf{r}}$ under the associated action map $\texttt{rep} \times K^\times \to \texttt{rep}$, and $\widetilde{\Omega}_{\mathbf{r}}$ as the closure of $\widetilde{\Omega}_{\mathbf{r}}^\circ$ in $\texttt{rep} \times \mathbb{A}^1$, where again we consider $K^\times \subset \mathbb{A}^1$. We use $\widetilde{\Omega}_{\mathbf{r}}(a)$ to denote the fiber over $a \in \mathbb{A}^1$.

Lemma 4.9. *The family $\widetilde{\Omega_{\mathbf{r}}} \subseteq$ `rep` $\times \mathbb{A}^1$ is flat over $\mathbb{A}^1$.*

Proof. Let f_{ij}^α be the coordinate function picking out the (i,j)-entry of a matrix of V_α in our fixed basis, and define f_{ij}^β similarly. The action of $K^\times$ on the coordinate ring of `rep` induced from the action above is

$$f_{ij}^\alpha \cdot t = t^{-(i+j)} f_{ij}^\alpha \qquad f_{ij}^\beta \cdot t = t^{i+j} f_{ij}^\beta. \tag{4.10}$$

Let $I_{\mathbf{r}} \subseteq K[\mathtt{rep}]$ be the ideal defining $\Omega_{\mathbf{r}}$. The defining ideal of $\widetilde{\Omega_{\mathbf{r}}}$ in $\mathtt{rep} \times \mathbb{A}^1$ is exactly the homogenization $\widetilde{I_{\mathbf{r}}} \subseteq K[\mathtt{rep} \times \mathbb{A}^1]$ of $I_{\mathbf{r}}$ with respect to the above $K^\times$-action (see §2.11), thus the family is flat by Theorem 2.52. ☐

The proof of the following lemma is similar to [KMS06, Lemma 4.10] and omitted.

Lemma 4.11. *If a lacing diagram* $\mathbf{w}$ *is in the fiber* $\widetilde{\Omega}_{\mathbf{r}}(a_0)$ *for some* $a_0 \in K^\times$*, then* $\mathbf{w}$ *is in* $\widetilde{\Omega}_{\mathbf{r}}(a)$ *for all* $a \in \mathbb{A}^1$*.*

Action of $\widetilde{GL}$ *on* $\widetilde{\mathtt{rep}}$*.* We take the following right action of GL^2 on $\mathtt{rep}$, again omitting arrow and vertex subscripts since there is only one possible interpretation of which vertex group with a given α or β label acts on each side of a map.

$$(V_\beta,\ V_\alpha) \cdot (g_x^\beta, g_x^\alpha, g_y^\alpha, g_y^\beta) = ((g_y^\beta)^{-1} V_\beta\ g_x^\beta,\ (g_y^\alpha)^{-1} V_\alpha\ g_x^\alpha)$$

The following diagrammatic illustration may be helpful.

$$\begin{tikzcd}
 & GL_{y_i}^{\alpha} \times GL_{y_i}^{\beta} & & \\
 & y_i \arrow[dl, "\mathtt{rep}(\alpha_{i+1})"'] \arrow[dr, "\mathtt{rep}(\beta_i)"] & & {} \arrow[dl, "\mathtt{rep}(\alpha_i)"] \\
{} & & x_i & \\
 & & GL_{x_i}^{\beta} \times GL_{x_i}^{\alpha} &
\end{tikzcd} \tag{4.12}$$

From equations (4.2), (4.3), and (4.8), we see that this action is $K^\times$-equivariant in the sense that

$$(V \cdot t) \cdot (g \cdot t) = (V \cdot g) \cdot t \qquad \text{for all } t \in K^\times,\ g \in GL^2,\ V \in \mathtt{rep}. \tag{4.13}$$

Proposition 4.14. *The family* $\widetilde{GL}$ *acts fiberwise on* $\widetilde{\Omega}_{\mathbf{r}}$*.*

Proof. Using [KMS06, Prop. 4.1], it is enough to prove the statement for the restriction to fibers over $K^\times$. The action of $K^\times$ gives an isomorphism

$$GL \times K^\times \to \widetilde{GL}(t \neq 0), \qquad (g, t) \mapsto (g \cdot t, t). \tag{4.15}$$

There is also an isomorphism of subschemes of $\mathtt{rep} \times K^\times$

$$\Omega_{\mathbf{r}} \times K^\times \to \widetilde{\Omega}_{\mathbf{r}}(t \neq 0), \qquad (V, t) \mapsto (V \cdot t, t). \tag{4.16}$$

Consider the standard (fiberwise) action of the left hand side of (4.15) on the left hand side of (4.16). Showing that this action is compatible with these isomorphisms is equivalent to showing that $(V \cdot t) \cdot (g \cdot t) = (V \cdot g) \cdot t$ for all $t \in K^\times$, which is exactly (4.13). ☐

The next proposition is proven just as in the equioriented case (cf. [KMS06, Prop. 3.4]). Recall that $\mathcal{O}(\mathbf{w})$ is the product of matrix Schubert varieties corresponding to the partial permutations constituting $\mathbf{w}$, as defined in (2.40).

Proposition 4.17. *The groups* $B_+ \times B_-$ *and* $B_+ \times_T B_-$ *have the same orbits on* $\mathtt{rep}$*. Furthermore, each orbit contains a unique lacing diagram, and the orbit closures are exactly* $\mathcal{O}(\mathbf{w})$ *as* $\mathbf{w}$ *varies over all lacing diagrams.* ☐

Corollary 4.18. *For any rank conditions* $\mathbf{r}$*, the induced reduced subscheme of the special fiber* $\widetilde{\Omega}_{\mathbf{r}}(0)$ *is a union of various* $\mathcal{O}(\mathbf{w})$*. Furthermore, since the family* $\widetilde{\Omega}_{\mathbf{r}}$ *is flat, each irreducible component of* $\widetilde{\Omega}_{\mathbf{r}}(0)$ *has the same dimension as* $\Omega_{\mathbf{r}}$*.*

Reducedness of the initial scheme. Our next goal is to show that $\widetilde{\Omega}_{\mathbf{r}}(0)$ is reduced. To do this, we consider a degeneration of the Zelevinsky image $\zeta(\Omega_{\mathbf{r}})$.

Let $d = d_x + d_y$ and recall that $Y_\circ^{v_0}$ denotes the set of $d \times d$ matrices of the form

$$\begin{bmatrix} * & \mathbf{1}_{d_y} \\ \mathbf{1}_{d_x} & 0 \end{bmatrix},$$

where the entries in the $*$ block are arbitrary elements of the base field K. Let Z be a matrix of this type, but which has indeterminates (instead of field elements) in the northwest block of $*$ entries. We denote this submatrix of variables by $Z_{d_y \times d_x}$. We may thus identify the coordinate ring $K[Y_\circ^{v_0}]$ with the polynomial ring $K[z_{ij}]$ generated by z_{ij}, $1 \leq i \leq d_y$, $1 \leq j \leq d_x$. We assign weights to the various z_{ij} as follows, recalling Definition 2.13 of the snake region.

1. (1) Suppose that z_{ij} lies in a block in the snake region of Z associated to a left-pointing, β-type arrow. If z_{ij} is the (k,l)-entry of its block, assign weight $\lambda(z_{ij}) = k + l$.
2. (2) Suppose that z_{ij} lies in a block in the snake region of Z associated to a right-pointing, α-type arrow. If z_{ij} is the (k,l)-entry of its block, assign weight $\lambda(z_{ij}) = -(k + l)$.
3. (3) The weights of the variables outside of the snake region will not matter, and so we assign weight $\lambda(z_{ij}) = 0$ to every z_{ij} outside of the snake region.

In Lemma 4.26 below, we will see that the integral weight function λ is constructed so that

$$\mathrm{in}_\lambda \zeta(\Omega_{\mathbf{r}}) \cong \widetilde{\Omega}_{\mathbf{r}}(0).$$

Consequently, reducedness of $\mathrm{in}_\lambda \zeta(\Omega_{\mathbf{r}})$, which is shown in Lemma 4.22, will imply reducedness of $\widetilde{\Omega}_{\mathbf{r}}(0)$.

Example 4.19. We continue Example 2.15. The matrix $Z_{d_y \times d_x}$ of indeterminates appears below on the left, in which the variables in the snake region have been renamed to make it easier to see the block structure. The corresponding weights of the variables appear below on the right.

$$\begin{bmatrix}
z_{1,1} & z_{1,2} & z_{1,3} & z_{1,4} & z_{1,5} & a_{1,1} & a_{1,2} \\
z_{2,1} & z_{2,2} & c_{1,1} & c_{1,2} & c_{1,3} & b_{1,1} & b_{1,2} \\
z_{3,1} & z_{3,2} & c_{2,1} & c_{2,2} & c_{2,3} & b_{2,1} & b_{2,2} \\
e_{1,1} & e_{1,2} & d_{1,1} & d_{1,2} & d_{1,3} & z_{4,6} & z_{4,7} \\
e_{2,1} & e_{2,2} & d_{2,1} & d_{2,2} & d_{2,3} & z_{5,6} & z_{5,7} \\
f_{1,1} & f_{1,2} & z_{6,3} & z_{6,4} & z_{6,5} & z_{6,6} & z_{6,7} \\
f_{1,2} & f_{2,2} & z_{7,3} & z_{7,4} & z_{7,5} & z_{7,6} & z_{7,7}
\end{bmatrix}, \quad
\begin{bmatrix}
0 & 0 & 0 & 0 & 0 & 2 & 3 \\
0 & 0 & 2 & 3 & 4 & -2 & -3 \\
0 & 0 & 3 & 4 & 5 & -3 & -4 \\
2 & 3 & -2 & -3 & -4 & 0 & 0 \\
3 & 4 & -3 & -4 & -5 & 0 & 0 \\
-2 & -3 & 0 & 0 & 0 & 0 & 0 \\
-3 & -4 & 0 & 0 & 0 & 0 & 0
\end{bmatrix} \qquad \square$$

We will also need the monomial order $\prec$ on $K[z_{ij}]$ from [WY12, §2.3]. This is defined as the columnwise lexicographic order down columns of $Z_{d_y \times d_x} = (z_{ij})$, starting in the northeast: $z_{ij} \prec z_{kl}$ when either $j < l$, or $j = l$ and $k < i$. Note that the lexicographically greatest variable in any submatrix of $Z_{d_y \times d_x}$ is always in the northeast corner. Observe then that $\prec$ is an *antidiagonal monomial order*, meaning that the leading term of any minor in Z is the product of the antidiagonal entries of the corresponding submatrix (the *antidiagonal term* of the minor).

Lemma 4.20. *Let Z' be an $n \times n$ submatrix of $Z_{d_y \times d_x}$ and let h denote the polynomial obtained from $\det(Z')$ by setting all variables outside of the snake region to zero. Then, for λ and $\prec$ as above,*

$$\mathrm{in}_\prec \mathrm{in}_\lambda h = \mathrm{in}_\prec h. \tag{4.21}$$

Furthermore, if h is not the zero polynomial, the above initial term is, up to sign, the antidiagonal term of Z'.

Proof. We first claim that if h is not the zero polynomial, then, up to sign, one of the terms of h is the antidiagonal term. Indeed, if h does not contain the antidiagonal term, then some antidiagonal entry of Z' is a variable outside of the snake region, which was set to 0. Thus Z' has an antidiagonal entry z_{ij} from outside of the snake region of $Z_{d_y \times d_x}$. By the block structure of $Z_{d_y \times d_x}$, one of the following holds: either all entries in Z' which are weakly southeast of z_{ij} are outside of the snake region of $Z_{d_y \times d_x}$, or all entries in Z' which are weakly northwest of z_{ij} are outside of the snake region of $Z_{d_y \times d_x}$. In either case, if the variables outside of the snake region are set to 0, there is no specialization of the other variables of Z' which results in a full rank matrix. Consequently, h is the zero polynomial.

Next observe that the antidiagonal term of h is a term of $\mathrm{in}_\lambda h$. To this end, note that the terms of h are indexed by $n \times n$ permutations in the usual way: up to sign, the term of h corresponding to $\sigma \in S_n$ is the product of those z_{ij} such that the permutation matrix σ^T has a 1 in location (i,j). Let h_σ denote the term associated to σ. Suppose σ_1^T and σ_2^T are two $n \times n$ permutation matrices which are identical except for the following: σ_1^T has ones in locations (i,j) and (k,l), with $i < k$, $j < l$, and σ_2^T has ones in locations (i,l) and (k,j) so that σ_2 has more inversions than σ_1. There are three possible cases if both σ_1 and σ_2 correspond to (nonzero) terms of h:

1. If locations (i,j) and (k,l) are in the same block of the snake, then $\lambda(h_{\sigma_1}) = \lambda(h_{\sigma_2})$.
2. If (i,j) and (k,l) are in adjacent horizontal blocks of the snake, then $\lambda(h_{\sigma_1}) < \lambda(h_{\sigma_2})$.
3. If (i,j) and (k,l) are in adjacent vertical blocks of the snake, then $\lambda(h_{\sigma_1}) < \lambda(h_{\sigma_2})$.

Thus, the antidiagonal term of h, which is indexed by the longest element $w_0 \in S_n$, must have maximal possible weight, and so is a term of $\mathrm{in}_\lambda h$. It now follows immediately from the definition of $\prec$ that the monomial $\mathrm{in}_\prec \mathrm{in}_\lambda h$ is the antidiagonal term of $\det(Z')$ (which is also $\mathrm{in}_\prec h$). ☐

Lemma 4.22. *Let $I_{v(\mathbf{r})} \subseteq K[z_{ij}]$ denote the defining ideal of the Zelevinsky image $\zeta(\Omega_\mathbf{r})$ in $Y_\circ^{v_0}$. For the weight function λ and the monomial order $\prec$ described above, we have*

$$in_\prec in_\lambda I_{v(\mathbf{r})} = in_\prec I_{v(\mathbf{r})}, \tag{4.23}$$

which is a squarefree monomial ideal. Consequently, $in_\lambda I_{v(\mathbf{r})}$ is a radical ideal.

Proof. Let $\mathcal{E}ss(v(\mathbf{r}))$ denote the essential set of the Zelevinsky permutation $v(\mathbf{r})$ (see §2.5 for the definition of essential set). Since $\zeta(\Omega_\mathbf{r})$ is a Kazhdan-Lusztig variety, we may apply [WY12, Theorem 2.1] to see that the set of essential minors

$$G_{v(\mathbf{r})} := \{\text{minors of size } (1 + \operatorname{rank} v(\mathbf{r})_{p\times q}) \text{ in } Z_{p\times q} \mid (p,q) \in \mathcal{E}ss(v(\mathbf{r}))\}$$

is a Gröbner basis of $I_{v(\mathbf{r})}$ for the monomial order $\prec$. Next observe that any non-constant minor of

$$Z = \begin{bmatrix} * & \mathbf{1}_{d_y} \\ \mathbf{1}_{d_x} & 0 \end{bmatrix} \tag{4.24}$$

is, up to sign, equal to a minor of the northwest $*$ block of variables, $Z_{d_y \times d_x}$. By Lemma 4.20, each such minor h has the property that $\mathrm{in}_\prec \mathrm{in}_\lambda h = \mathrm{in}_\prec h$, so we have

$$\mathrm{in}_\prec I_{v(\mathbf{r})} \subseteq \mathrm{in}_\prec \mathrm{in}_\lambda I_{v(\mathbf{r})}. \tag{4.25}$$

Both ideals in (4.25) are initial ideals of $I_{v(\mathbf{r})} \subseteq K[z_{ij}]$, so their associated quotient rings both have the same Hilbert series as $K[z_{ij}]/I_{v(\mathbf{r})}$ by [MS05, Prop. 8.28] and the translation between K-polynomials and Hilbert series. Thus, the Hilbert series of these two rings are equal. Then the containment (4.25) implies that the two ideals $\mathrm{in}_\prec I_{v(\mathbf{r})}$ and $\mathrm{in}_\prec \mathrm{in}_\lambda I_{v(\mathbf{r})}$ are equal. Finally, the initial ideal $\mathrm{in}_\prec \mathrm{in}_\lambda I_{v(\mathbf{r})}$ is radical since it is a squarefree monomial ideal, so the ideal $\mathrm{in}_\lambda I_{v(\mathbf{r})}$ is also radical. ☐

Finally, we observe the following:

Lemma 4.26. *The degeneration* $\Omega_{\mathbf{r}} \rightsquigarrow \widetilde{\Omega}_{\mathbf{r}}(0)$ *is compatible with the Zelevinsky map* ζ, *meaning that*

$$\operatorname{Spec} K[z_{ij}]/in_\lambda I_{v(\mathbf{r})} \cong \widetilde{\Omega}_{\mathbf{r}}(0).$$

Consequently, $\widetilde{\Omega}_{\mathbf{r}}(0)$ *is reduced.*

Proof. Let $\zeta^\sharp : K[z_{ij}] \to K[\mathtt{rep}]$ be the surjective ring homomorphism induced by the Zelevinsky map $\zeta\colon \mathtt{rep} \to Y_\circ^{v_0}$. Then, $\zeta^\sharp$ maps z_{ij} to 0 if (i,j) is a position outside of the snake region, and $\zeta^\sharp$ maps each snake region variable z_{ij} to its associated $f_{kl}^{\alpha_m}$ or $f_{kl}^{\beta_m}$ in $K[\mathtt{rep}]$. In this proof, we identify each $f_{kl}^{\alpha_m}$ (or $f_{kl}^{\beta_m}$) with its associated snake region variable z_{ij} in $K[z_{ij}]$; it will be clear from the context whether a given polynomial involving only snake region variables is an element of $K[z_{ij}]$ or of $K[\mathtt{rep}]$.

Let $\widetilde{I}_{\mathbf{r}}(0)$ denote the scheme-theoretic defining ideal of $\widetilde{\Omega}_{\mathbf{r}}(0)$. We need to check that

$$(\zeta^\sharp)^{-1}(\widetilde{I}_{\mathbf{r}}(0)) = \mathrm{in}_\lambda I_{v(\mathbf{r})}.$$

To do so, suppose that $\{g_1, \dots, g_k\}$ is a Gröbner basis of the quiver ideal $I_{\mathbf{r}} \subseteq K[\mathtt{rep}]$ for the integral weight function μ determined by the $K^\times$-action described in (4.10). Then, the set

$$\{g_1, \dots, g_k\} \cup \{z_{ij} \mid (i,j) \text{ not in snake region of } M\} \tag{4.27}$$

generates the ideal $(\zeta^\sharp)^{-1}(I_{\mathbf{r}}) = I_{v(\mathbf{r})}$. Since each polynomial g_i involves only snake region variables, the set (4.27) is a Gröbner basis of $I_{v(\mathbf{r})}$ for the integral weight function λ because the integral weight functions μ and λ assign the same weights to snake region variables. Consequently,

$$\begin{aligned}
\mathrm{in}_\lambda I_{v(\mathbf{r})} &= \langle \mathrm{in}_\lambda g_i \mid 1 \le i \le k\rangle + \langle z_{ij} \mid (i,j) \text{ not in snake region}\rangle \\
&= \langle \mathrm{in}_\mu g_i \mid 1 \le i \le k\rangle + \langle z_{ij} \mid (i,j) \text{ not in snake region}\rangle \\
&= \mathrm{in}_\mu I_{\mathbf{r}} + \langle z_{ij} \mid (i,j) \text{ not in snake region}\rangle \\
&= \widetilde{I}_{\mathbf{r}}(0) + \langle z_{ij} \mid (i,j) \text{ not in snake region}\rangle \\
&= (\zeta^\sharp)^{-1}(\widetilde{I}_{\mathbf{r}}(0))
\end{aligned}$$

as desired. It now follows from Lemma 4.22 that $\widetilde{I}_{\mathbf{r}}(0)$ is radical. □

We are now ready to prove the main theorem of this section.

Theorem 4.28. *Any type A quiver locus degenerates to a reduced union of products of matrix Schubert varieties. More precisely, for a quiver locus* $\Omega_{\mathbf{r}} \subseteq \mathtt{rep}$ *we have a degeneration*

$$\widetilde{\Omega}_{\mathbf{r}}(0) = \bigcup_{\mathbf{w} \in W(\mathbf{r})} \mathcal{O}(\mathbf{w}), \tag{4.29}$$

where the union is scheme-theoretic and taken over minimal lacing diagrams for $\mathbf{r}$.

Proof. By Lemma 4.26, $\widetilde{\Omega}_{\mathbf{r}}(0)$ is a reduced scheme. So, by Corollary 4.18, the irreducible components of $\widetilde{\Omega}_{\mathbf{r}}(0)$ are a collection of $\mathcal{O}(\mathbf{w})$. Let J denote the set of lacing diagrams $\mathbf{w}$ where $\mathcal{O}(\mathbf{w})$ is an irreducible component of $\widetilde{\Omega}_{\mathbf{r}}(0)$. We will show that J is the set $W(\mathbf{r})$ of minimal lacing diagrams for $\mathbf{r}$.

By Lemma 4.11, we see that for each lacing diagram $\mathbf{w} \in \Omega_{\mathbf{r}} = \widetilde{\Omega}_{\mathbf{r}}(1)$, we have that $\mathbf{w} \in \widetilde{\Omega}_{\mathbf{r}}(0)$. Now suppose that $\mathbf{w}$ is a minimal lacing diagram for $\mathbf{r}$, so that the number of its crossings is the codimension of $\Omega_{\mathbf{r}}$ in $\mathtt{rep}$. The orbit closure of $\mathbf{w}$ in the the fiber over $t = 0$ is its $B_+ \times_T B_-$ orbit closure $\mathcal{O}(\mathbf{w})$, whose codimension in $\mathtt{rep}$ is also counted by the number of its crossings. Thus the orbit closure of $\mathbf{w}$ in $\widetilde{\Omega}_{\mathbf{r}}(0)$ is of maximal possible dimension and an irreducible component. Therefore, $\widetilde{\Omega}_{\mathbf{r}}(0)$ contains the union of all $\mathcal{O}(\mathbf{w})$ as $\mathbf{w}$ runs over the set of minimal lacing diagrams for $\mathbf{r}$, that is, $J \supseteq W(\mathbf{r})$.

We now show that there are no other irreducible components of $\widetilde{\Omega}_{\mathbf{r}}(0)$, and thus that $J \subseteq W(\mathbf{r})$. Consider the sequence of degenerations

$$\Omega_{\mathbf{r}} \rightsquigarrow \widetilde{\Omega}_{\mathbf{r}}(0) = \bigcup_{\mathbf{w}\in J} \mathcal{O}(\mathbf{w}) \rightsquigarrow \mathrm{in}_{\prec}\left(\bigcup_{\mathbf{w}\in J} \mathcal{O}(\mathbf{w})\right) = \bigcup_{\mathbf{w}\in J} \mathrm{in}_{\prec}\mathcal{O}(\mathbf{w}). \tag{4.30}$$

The rightmost equality in (4.30) holds since $\mathrm{in}_{\prec}\left(\bigcup_{\mathbf{w}\in J} \mathcal{O}(\mathbf{w})\right)$ is reduced (see Lemma 4.22).

Let $P \in \mathrm{Pipes}(v_0, v(\mathbf{r}))$ and let $H_P \subseteq Y_{\circ}^{v_0}$ denote the coordinate subspace with defining ideal

$$I_P := \langle z_{ij} \mid + \in P \text{ is in location } (i,j)\rangle \subseteq K[z_{ij}].$$

Recall that each $\mathcal{O}(\mathbf{w})$ is a product of matrix Schubert varieties, where the factors are indexed by the blocks in the snake region of the Zelevinsky image. Thus,

$$\bigcup_{\mathbf{w}\in J} \mathrm{in}_{\prec}\mathcal{O}(\mathbf{w}) = \bigcup_{\mathbf{w}\in J}\bigcup_{P} H_P \tag{4.31}$$

where the rightmost union in (4.31) is taken over those pipe dreams P which are reduced within each block of the snake region, and which have the property $\mathbf{w}(P) = \mathbf{w}$ (see §2.10).

By Lemma 4.22, the sequence of two degenerations in (4.30) results in the scheme $\mathrm{in}_{\prec}\Omega_{\mathbf{r}}$. By [WY12, Theorem 3.2],

$$\mathrm{in}_{\prec}\Omega_{\mathbf{r}} = \bigcup_{P\in \mathrm{RedPipes}(v_0, v(\mathbf{r}))} H_P. \tag{4.32}$$

Together (4.31) and (4.32), imply that for each irreducible component $\mathcal{O}(\mathbf{w})$ of $\widetilde{\Omega}_{\mathbf{r}}(0)$, there is some $P \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))$ such that $\mathbf{w}(P) = \mathbf{w}$. By Proposition 2.50, we know that $\mathbf{w} = \mathbf{w}(P)$ is an element of the minimal lacing diagrams $W(\mathbf{r})$. Thus, $J \subseteq W(\mathbf{r})$. ☐

4.2. **Component formulas.**

For a lacing diagram $\mathbf{w}$ associated to a bipartite type A quiver, we denote the constituent sequence of partial permutation matrices as

$$\mathbf{w} = (w_n, \widetilde{w}_n, \ldots, w_2, \widetilde{w}_2, w_1, \widetilde{w}_1)$$

where w_i is the matrix associated to β_i and $\widetilde{w}_i$ to α_i. Recall the completion $c(w_i)$ and opposite completion $oc(\widetilde{w}_i)$ of partial permutation matrices from §2.5. To a lacing diagram, we assign the following products of double Grothendieck and Schubert polynomials as a shorthand:

$$\mathfrak{G}_{\mathbf{w}}(\mathbf{t};\mathbf{s}) = \mathfrak{G}_{c(w_n)}(\mathbf{t}^n;\mathbf{s}^n)\mathfrak{G}_{oc(\widetilde{w}_n)}(\widetilde{\mathbf{t}}^{n-1};\widetilde{\mathbf{s}}^n)\cdots\mathfrak{G}_{c(w_1)}(\mathbf{t}^1;\mathbf{s}^1)\mathfrak{G}_{oc(\widetilde{w}_1)}(\widetilde{\mathbf{t}}^0;\widetilde{\mathbf{s}}^1) \tag{4.33}$$

$$\mathfrak{S}_{\mathbf{w}}(\mathbf{t};\mathbf{s}) = \mathfrak{S}_{c(w_n)}(\mathbf{t}^n;\mathbf{s}^n)\mathfrak{S}_{oc(\widetilde{w}_n)}(\widetilde{\mathbf{t}}^{n-1};\widetilde{\mathbf{s}}^n)\cdots\mathfrak{S}_{c(w_1)}(\mathbf{t}^1;\mathbf{s}^1)\mathfrak{S}_{oc(\widetilde{w}_1)}(\widetilde{\mathbf{t}}^0;\widetilde{\mathbf{s}}^1). \tag{4.34}$$

In this notation, we have by (2.27) and (2.40) that the K-polynomial of $\mathcal{O}(\mathbf{w})$ with respect to its embedding in $\mathtt{rep}(\mathbf{d})$ is $\mathfrak{G}_{\mathbf{w}}(\mathbf{t};\mathbf{s})$.

Recall that the Möbius function of a finite poset is the (unique) function $\mu\colon P \to \mathbb{Z}$ such that $\sum_{y\geq x}\mu_P(y) = 1$ for all $x \in P$. It is typically used in the following way: for $x \in P$, we define "open" and "closed" characteristic functions

$$\chi_x^{\circ}(y) = \begin{cases}1 & y = x\\ 0 & y \neq x,\end{cases} \qquad \chi_x(y) = \begin{cases}1 & y \geq x\\ 0 & y \not\geq x.\end{cases} \tag{4.35}$$

Then, by inclusion-exclusion, we have that the constant function taking value 1 on P can be written as

$$1_P = \sum_x \chi_x^\circ = \sum_x \mu_P(x)\chi_x. \tag{4.36}$$

Theorem 4.37 (Bipartite component formula)**.** *Let* $\mathbf{r}$ *be a rank array for a bipartite, type* A *quiver* Q*. Then we have*

$$K\mathcal{Q}_{\mathbf{r}}(\mathbf{t}/\mathbf{s}) = \sum_{\mathbf{w}\in KW(\mathbf{r})} (-1)^{|\mathbf{w}|-\ell(v(\mathbf{r}))}\mathfrak{G}_{\mathbf{w}}(\mathbf{t};\mathbf{s}) \tag{4.38}$$

and

$$\mathcal{Q}_{\mathbf{r}}(\mathbf{t}-\mathbf{s}) = \sum_{\mathbf{w}\in W(\mathbf{r})} \mathfrak{S}_{\mathbf{w}}(\mathbf{t};\mathbf{s}) \tag{4.39}$$

Proof. By Theorem 4.28 and the fact that degeneration also preserves K-polynomials [MS05, Thereom 8.36], we have that

$$K\mathcal{Q}_{\mathbf{r}}(\mathbf{t}/\mathbf{s}) = \mathcal{K}_{\mathtt{rep}}(\widetilde{\Omega}_{\mathbf{r}}(0);\mathbf{s},\mathbf{t}) \tag{4.40}$$

Define the subset of lacing diagrams

$$\mathcal{M} := \{\mathbf{w} \mid \mathcal{O}(\mathbf{w}) \subseteq \widetilde{\Omega}_{\mathbf{r}}(0)\}, \tag{4.41}$$

which is partially ordered by coordinate-wise containment (i.e. $\mathbf{w} \leq \mathbf{w}'$ if and only if $\mathcal{O}(\mathbf{w}) \supseteq \mathcal{O}(\mathbf{w}')$). The collection of closed subvarieties $\{\mathcal{O}(\mathbf{w}) \subseteq \widetilde{\Omega}_{\mathbf{r}}(0) \mid \mathbf{w} \in \mathcal{M}\}$ has the intersect-decompose property of [Knu] because the intersection of any collection of matrix Schubert varieties is a union of matrix Schubert varieties. The hypotheses of [Knu, Theorem 1] are satisfied by this collection because Schubert varieties in a flag variety are simultaneously compatibly F-split [MR85], so we have that

$$\mathcal{K}_{\mathtt{rep}}(\widetilde{\Omega}_{\mathbf{r}}(0);\mathbf{s},\mathbf{t}) = \sum_{\mathbf{w}\in\mathcal{M}} \mu_{\mathcal{M}}(\mathbf{w})[\mathcal{O}(\mathbf{w})], \tag{4.42}$$

where we introduce the temporary shorthand $[\mathcal{O}(\mathbf{w})] = \mathcal{K}_{\mathtt{rep}}(\mathcal{O}(\mathbf{w});\mathbf{s},\mathbf{t})$ for this proof.

This Möbius function may be difficult to compute, so we pass to a poset where we can compute the Möbius function: the pipe complex of [WY12, §3]. Let $\Delta := \Delta_{v_0,v(\mathbf{r})}$ be the set of pipe dreams on the diagram of v_0 which contain a reduced pipe dream for $v(\mathbf{r})$. The partial order on Δ is determined by reverse containment of +s. We can consider Δ as a simplicial complex in this way, whose maximal faces are the elements of $\mathrm{RedPipes}(v_0, v(\mathbf{r}))$. It is known that Δ is always homeomorphic to a ball, since there is a unique pipe dream for v_0, and that the Möbius function is

$$\mu_\Delta(P) = \begin{cases} (-1)^{\operatorname{codim}_\Delta P} & \text{if } \delta(P) = v(\mathbf{r}) \\ 0 & \text{otherwise,} \end{cases} \tag{4.43}$$

where $\operatorname{codim}_\Delta P = |P| - \ell(v(\mathbf{r}))$ ([WY12, Prop. 3.3] via [KM04, Thm. 3.7]).

Since every element of Δ contains a pipe dream for $v(\mathbf{r})$, every element contains all +s outside the snake, so we can restrict our attention to the blocks of the snake as usual. We define an order-preserving map of posets $\pi\colon \Delta \to \mathcal{M}$ by $\pi(P) = \mathbf{w}(P)$, where $\mathbf{w}(P)$ is the lacing diagram associated to P in §2.10. To see that π does actually take values in $\mathcal{M}$, we need to show that $\mathbf{w}(P) \in \mathcal{M}$ whenever $P \in \Delta$. This is true when P is a reduced pipe dream for $v(\mathbf{r})$ because then $\mathbf{w}(P)$ is a minimal lacing diagram by Proposition 2.50 (a), so $\mathbf{w}(P)$ lies in $\widetilde{\Omega}_{\mathbf{r}}(0)$ by Theorem 4.28. For arbitrary $P \in \Delta$, choose a $P_0 \in \mathrm{RedPipes}(v_0, v(\mathbf{r}))$ such that $P_0 \subseteq P$, so that $P \leq P_0$ in Δ and thus $\mathbf{w}(P) \leq \mathbf{w}(P_0) \in \mathcal{M}$, so $\mathbf{w}(P) \in \mathcal{M}$ as well.

This induces a map on rings of $\mathbb{Z}$-valued functions

$$\pi^*\colon \mathbb{Z}[\mathcal{M}] \to \mathbb{Z}[\Delta].$$

By (4.36), in $\mathbb{Z}[\Delta]$ we have that

$$1_\Delta = \sum_{P\in\Delta} \chi_P^\circ = \sum_{P\in \mathrm{Pipes}(v_0,v(\mathbf{r}))} (-1)^{\operatorname{codim}_\Delta P}\chi_P, \tag{4.44}$$

where the last equality uses (4.43). Similarly, in $\mathbb{Z}[\mathcal{M}]$ we have the expression

$$1_\mathcal{M} = \sum_{\mathbf{w}\in\mathcal{M}} \chi_\mathbf{w}^\circ = \sum_{\mathbf{w}\in\mathcal{M}} \mu_\mathcal{M}(\mathbf{w})\chi_\mathbf{w} \tag{4.45}$$

which gives another way of writing

$$1_\Delta = \pi^*(1_\mathcal{M}) = \sum_{\mathbf{w}\in\mathcal{M}} \mu_\mathcal{M}(\mathbf{w})\pi^*(\chi_\mathbf{w}). \tag{4.46}$$

To compare this with (4.44), we need to express $\pi^*(\chi_\mathbf{w})$ in terms of the basis $\{\chi_P\}$. The claim is that

$$\pi^*(\chi_\mathbf{w}) = \sum_{\substack{P\in\Delta\\ \pi(P)=\mathbf{w}}} (-1)^{\operatorname{codim}_{\Delta(\mathbf{w})} P}\chi_P, \tag{4.47}$$

where $\Delta(\mathbf{w}) := \{P\in\Delta \mid \pi(P)\geq \mathbf{w}\}$ and $\operatorname{codim}_{\Delta(\mathbf{w})} P = |P| - |\mathbf{w}|$. Now notice that $\Delta(\mathbf{w})$ is just a product of pipe complexes

$$\Delta(\mathbf{w}) = \prod_{i=1}^{n} \Delta_{\alpha_i}(\widetilde{w}_i)\times\Delta_{\beta_i}(w_i) \tag{4.48}$$

where $\Delta_{\alpha_i}(\widetilde{w}_i)$ is the complex of pipe dreams on the α_i-block of the snake which contain a pipe dream for $\widetilde{w}_i$, and similarly for $\Delta_{\beta_i}(w_i)$. Thus, $\Delta(\mathbf{w})$ is also homeomorphic to a ball and its Möbius function can be computed in each factor in the same way as in (4.43). We get

$$\mu_{\Delta(\mathbf{w})}(P) = \begin{cases} (-1)^{\operatorname{codim}_{\Delta(\mathbf{w})} P} & \pi(P) = \mathbf{w}\\ 0 & \pi(P)\neq \mathbf{w}.\end{cases} \tag{4.49}$$

Now to verify the claim, we use the definition of $\pi^*(\chi_\mathbf{w})$ and Möbius inversion to compute

$$\pi^*(\chi_\mathbf{w}) = \sum_{\substack{P\in\Delta\\ \pi(P)\geq\mathbf{w}}} \chi_P^\circ = \sum_{\substack{P\in\Delta\\ \pi(P)=\mathbf{w}}} (-1)^{\operatorname{codim}_{\Delta(\mathbf{w})} P}\chi_P. \tag{4.50}$$

Substituting this into (4.46) we get

$$1_\Delta = \sum_{\mathbf{w}\in\mathcal{M}} \mu_\mathcal{M}(\mathbf{w}) \left(\sum_{\substack{P\in\Delta\\ \pi(P)=\mathbf{w}}} (-1)^{\operatorname{codim}_{\Delta(\mathbf{w})} P}\chi_P \right) = \sum_{P\in\Delta} \mu_\mathcal{M}(\pi(P))(-1)^{\operatorname{codim}_{\Delta(\pi(P))} P}\chi_P \tag{4.51}$$

Comparing the coefficient of χ_P in (4.44) and (4.51) we find that

$$\mu_\mathcal{M}(\pi(P)) = \begin{cases} (-1)^{|\mathbf{w}|-\ell(v(\mathbf{r}))} & \text{if } P\in \mathrm{Pipes}(v_0, v(\mathbf{r}))\\ 0 & \text{otherwise}\end{cases} \tag{4.52}$$

Here, we have used that $\operatorname{codim}_\Delta P - \operatorname{codim}_{\Delta(\mathbf{w})} P = |\mathbf{w}| - \ell(v(\mathbf{r}))$, which just follows from the definitions.

Next we want to see that π is surjective. By Proposition 2.50, every minimal lacing diagram for $v(\mathbf{r})$ is in the image of π. For arbitrary $\mathbf{w} \in \mathcal{M}$, choose a minimal lacing diagram $\mathbf{w}_0 \geq \mathbf{w}$ (which exists by Theorem 4.28), and a lift $P_0 \in \operatorname{RedPipes}(v_0, v(\mathbf{r}))$ such that $\mathbf{w}(P_0) = \mathbf{w}_0$, and then in each block of the snake add +s to produce a pipe dream P such that $\mathbf{w}(P) = \mathbf{w}$. This is possible block-by-block because for each arrow of Q, the condition $\mathbf{w}_0 \geq \mathbf{w}$ means exactly that we have containment of matrix Schubert varieties indexed by this factor in $\mathcal{O}(\mathbf{w}_0) \supseteq \mathcal{O}(\mathbf{w})$. We have that $P \in \Delta$ since it contains a reduced pipe dream for $v(\mathbf{r})$ on v_0 by definition, so we have shown that π is surjective. Now this implies that every $\mathbf{w}$ appearing in the sum (4.42) is in the image of π, so by (4.52) the sum simplifies to a sum over $\mathbf{w}$ such that there exist $P \in \operatorname{Pipes}(v_0, v(\mathbf{r}))$ with $\pi(P) = \mathbf{w}$. By results of Buch, Fehér, and Rimányi ([Buc05, Lemma 6.2] and [BFR05, Theorem 3]), this is equivalent to summing over all K-theoretic lacing diagrams for $\mathbf{r}$.[1] Then the K-theoretic component formula follows immediately from (4.42) by substitutions from (4.40), (4.52), and (2.27). The multidegree version (4.39) is obtained from the K-polynomial formula as usual. □

5. GENERALIZING THE FORMULAS TO ARBITRARY ORIENTATION

5.1. **Reduction to the bipartite setting.** In this section, we recall the connection between quiver loci of arbitrarily oriented type A quivers and quiver loci in the bipartite setting. Utilizing this connection, we see that a simple substitution into our bipartite formulas yield the analogous formulas in the arbitrarily oriented setting. We begin by reviewing the relevant material from [KR15]. Since we work with more than one quiver and dimension vector, these are reinstated to the notation.

Example 5.1. Let Q be the quiver:

$$\begin{tikzcd}
 & z_4 \arrow[dl, "\gamma_5"'] \arrow[dr, "\gamma_4"] & & & & z_0 \arrow[dl, "\gamma_1"] \\
z_5 & & z_3 \arrow[dr, "\gamma_3"'] & & z_1 \arrow[dl, "\gamma_2"] & \\
 & & & z_2 & &
\end{tikzcd} \tag{5.2}$$

We construct an associated bipartite quiver $\widetilde{Q}$ by adding two new vertices w_1, w_3, and two new arrows δ_1, δ_3.

$$\begin{tikzcd}
 & z_4 \arrow[dl, "\gamma_5"'] \arrow[dr, "\gamma_4"] & & z_3 \arrow[dl, red, dashed, "\delta_3"] \arrow[dr, "\gamma_3"] & & \color{red}{w_1} \arrow[dl, "\gamma_2"'] \arrow[dr, red, dashed, "\delta_1"] & & z_0 \arrow[dl, "\gamma_1"] \\
z_5 & & \color{red}{w_3} & & z_2 & & z_1 &
\end{tikzcd}. \tag{5.3}$$

□

In general, we may define an associated bipartite quiver $\widetilde{Q}$ by the following local insertions of vertices and arrows: for each intermediate vertex of the form $z_{i+1} \xrightarrow{\gamma_{i+1}} z_i \xrightarrow{\gamma_i}$, add a sink w_i and arrow δ_i in the configuration

$$\begin{tikzcd}
z_{i+1} \arrow[dr, "\gamma_{i+1}"'] & & z_i \arrow[dl, "\delta_i"] \\
 & w_i &
\end{tikzcd}. \tag{5.4}$$

---

[1] Their work is in the equioriented case but readily generalizes to arbitrary orientation.

For each intermediate vertex of the form $z_{i+1} \xleftarrow{\gamma_{i+1}} z_i \xleftarrow{\gamma_i}$, add a source w_i and arrow δ_i in the configuration

$$\begin{tikzcd} & w_i \arrow[dl, "\gamma_{i+1}"'] \arrow[dr, "\delta_i"] & \\ z_{i+1} & & z_i \end{tikzcd} \tag{5.5}$$

The arrows and vertices coming from Q will be referred to as *original* arrows and vertices, and the new ones as *added* arrows and vertices. Additionally, the arrows δ_i will be sometimes referred to as *inverted arrows*, and anything indexed by them (blocks of the snake region, for example) can be referred to as "inverted" for reasons we will see below.

For a dimension vector $\mathbf{d}$ for Q, define $\widetilde{\mathbf{d}}$ as the natural lifting

$$\widetilde{\mathbf{d}}(z_i) := \mathbf{d}(z_i) \text{ and } \widetilde{\mathbf{d}}(w_i) := \mathbf{d}(z_i).$$

Let $G^* = \prod GL_{\widetilde{\mathbf{d}}(w_i)}(K)$ be the base change group at the added vertices, so that

$$GL(\widetilde{\mathbf{d}}) = G^* \times GL(\mathbf{d}). \tag{5.6}$$

Throughout this section, we denote a typical element of $\mathtt{rep}_{\widetilde{Q}}(\widetilde{\mathbf{d}})$ by $\widetilde{V} = (V_{\delta_i}) \times (V_{\gamma_i})$, and an element of $GL(\widetilde{\mathbf{d}})$ by $\widetilde{g} = (g_{w_i}) \times (g_{z_i})$. We write $U \subset \mathtt{rep}_{\widetilde{Q}}(\widetilde{\mathbf{d}})$ for the $GL(\widetilde{\mathbf{d}})$-stable open subscheme of representations for which every V_{δ_i} is invertible.

The following theorem collects the main facts from [KR15, Proposition 5.2 and Theorem 5.3] that we need.

Theorem 5.7. *Let Q be a quiver of type A, and $\widetilde{Q}$ the associated bipartite quiver defined above. Then there is a morphism*

$$\pi \colon U \to \mathtt{rep}_Q(\mathbf{d}) \tag{5.8}$$

which is equivariant with respect to the natural projection of base change groups

$$GL(\widetilde{\mathbf{d}}) \to GL(\mathbf{d}),$$

and also a principal G^*-bundle. Furthermore, each orbit closure $\overline{\mathcal{O}} \subseteq \mathtt{rep}_Q(\mathbf{d})$ for an arbitrary type A quiver is isomorphic to an open subset of an orbit closure of $\mathtt{rep}_{\widetilde{Q}}(\widetilde{\mathbf{d}})$, up to a smooth factor. Namely, we have

$$\overline{\pi^{-1}(\mathcal{O})} \simeq G^* \times \overline{\mathcal{O}}, \tag{5.9}$$

where the closure on the left hand side is taken in U.

The map π in the above theorem is defined as follows: for $\widetilde{V} \in U$, define matrices $X_{\gamma_i} = V_{\gamma_i}$ when γ_i is an arrow between two original vertices. We set $X_{\gamma_i} = V_{\gamma_i} V_{\delta_{i-1}}^{-1}$ or $X_{\gamma_i} = V_{\delta_{i-1}}^{-1} V_{\gamma_i}$ when γ_i is involved in a local configuration of type (5.4) or (5.5), respectively. Then define the projection map by

$$\pi \colon U \to \mathtt{rep}_Q(\mathbf{d}), \qquad \widetilde{V} \mapsto (X_{\gamma_i}). \tag{5.10}$$

Now, let σ denote the section of the map π which maps $(V_{\gamma_i}) \in \mathtt{rep}_Q(\mathbf{d})$ to $\widetilde{V} = (I_{\delta_i}) \times (V_{\gamma_i})$, where I_{δ_i} denotes the identity map over the arrow δ_i. Observe that σ is equivariant with respect to "diagonal" inclusion

$$GL(\mathbf{d}) \to G^* \times GL(\mathbf{d}), \qquad (g_{z_i}) \mapsto (g_{z_i}) \times (g_{z_i}). \tag{5.11}$$

That is, to each added vertex w_i, the inclusion assigns the group element g_{z_i} from the corresponding original vertex. This equivariance implies that the induced map on coordinate rings will respect the multigradings.

At this point we extend our notation for the multigrading of $K[\mathtt{rep}_Q(\mathbf{d})]$ beyond the bipartite orientation. The idea is to enlarge Q to the bipartite quiver $\widetilde{Q}$, take the alphabet we have already defined in that setting, then assign to each vertex of Q the alphabet of its corresponding vertex in $\widetilde{Q}$. To work with this convention explicitly, it is necessary to use simultaneously two notations for the arrows and vertices of $\widetilde{Q}$: the x, y, α, β labeling from previous sections and the z, w, γ, δ labeling indicating how it was obtained from Q. More precisely, for each vertex z_i of Q, we associate the alphabet

$$\begin{cases} \mathbf{t}^j & \text{if the associated vertex in } \widetilde{Q} \text{ is labeled } x_j \\ \mathbf{s}^j & \text{if the associated vertex in } \widetilde{Q} \text{ is labeled } y_j, \end{cases} \tag{5.12}$$

and then assign a degree to each coordinate function of each matrix in the same way as §2.6.

Let $\widetilde{d}$ be the total dimension of $\widetilde{\mathbf{d}}$ and d the total dimension of $\mathbf{d}$. As before, we can identify the multigrading groups of $K[\mathtt{rep}_{\widetilde{Q}}(\widetilde{\mathbf{d}})]$ and $K[\mathtt{rep}_Q(\mathbf{d})]$ with free abelian groups $\mathbb{Z}^{\widetilde{d}}$ and $\mathbb{Z}^d$, respectively, on the alphabets associated to the vertices. We define a quotient morphism of abelian groups $sub : \mathbb{Z}^{\widetilde{d}} \to \mathbb{Z}^d$ by identifying pairs alphabets for $\widetilde{Q}$ corresponding to the same vertex of Q. More precisely, for each original vertex of $\widetilde{Q}$, the corresponding alphabet is associated to the same vertex of Q, and sub acts as the identity on the corresponding group elements. For an added vertex of $\widetilde{Q}$ labeled x_i, for each $s^i_j \in \mathbf{s}^i$ we set $sub(s^i_j) = t^{i-1}_j$. For an added vertex of $\widetilde{Q}$ labeled y_i, for each $t^i_j \in \mathbf{t}^i$ we set $sub(t^i_j) = s^i_j$. In Figure 5 we have labeled the block rows and columns of the image of $\zeta \circ \sigma$ to show how sub interacts naturally with the Zelevinsky map, in the situation of Example 5.1.

$$\zeta(\sigma(V)) = \begin{array}{c|cccc|cccc|}
 & \mathbf{s}^3 & \mathbf{t}^2 & \mathbf{s}^2 & \mathbf{s}^1 & \mathbf{t}^0 & \mathbf{s}^1 & \mathbf{t}^2 & \mathbf{t}^3 \\
\mathbf{t}^0 & 0 & 0 & 0 & V_{\gamma_1} & \mathbf{1}_{\mathbf{d}(z_0)} & & & \\
\mathbf{s}^1 & 0 & 0 & V_{\gamma_2} & \mathbf{1}_{\mathbf{d}(z_1)} & & \mathbf{1}_{\mathbf{d}(z_1)} & & \\
\mathbf{t}^2 & 0 & \mathbf{1}_{\mathbf{d}(z_3)} & V_{\gamma_3} & 0 & & & \mathbf{1}_{\mathbf{d}(z_3)} & \\
\mathbf{t}^3 & V_{\gamma_5} & V_{\gamma_4} & 0 & 0 & & & & \mathbf{1}_{\mathbf{d}(z_4)} \\
\hline
\mathbf{s}^3 & \mathbf{1}_{\mathbf{d}(z_5)} & & & & & & & \\
\mathbf{t}^2 & & \mathbf{1}_{\mathbf{d}(z_3)} & & & & & & \\
\mathbf{s}^2 & & & \mathbf{1}_{\mathbf{d}(z_2)} & & & & 0 & \\
\mathbf{s}^1 & & & & \mathbf{1}_{\mathbf{d}(z_1)} & & & &
\end{array}$$

FIGURE 5. Example of the image of $\zeta \circ \sigma$

These definitions are made so that the restriction map on coordinate rings

$$\sigma^{\#} \colon K[\mathtt{rep}_{\widetilde{Q}}(\widetilde{\mathbf{d}})] \to K[\mathtt{rep}_Q(\mathbf{d})] \tag{5.13}$$

is graded with respect to the morphism sub of the multigrading groups. To simplify notation, we use the shorthand $\mathbf{s}, \mathbf{t}$ for the variable set of K-polynomials of objects associated to both Q and $\widetilde{Q}$, though strictly speaking one is a subset of the other. The map sub induces a map between the rings of Laurent polynomials associated to these alphabets, which we also denoted by sub.

Proposition 5.14. *The K-polynomial of an orbit closure for Q is obtained from the K-polynomial of its corresponding orbit closure for $\widetilde{Q}$ by substitution. That is, for an orbit closure $\overline{\mathcal{O}} \subseteq \mathtt{rep}_Q(\mathbf{d})$, we have $\mathcal{K}_{\mathtt{rep}_Q(\mathbf{d})}(\overline{\mathcal{O}};\mathbf{s},\mathbf{t}) = sub(\mathcal{K}_{\mathtt{rep}_{\widetilde{Q}}(\widetilde{\mathbf{d}})}(\overline{\pi^{-1}(\mathcal{O})};\mathbf{s},\mathbf{t}))$.*

Proof. Taking a $\mathbb{Z}^{\widetilde{d}}$-graded free resolution of $K[\overline{\pi^{-1}(\mathcal{O})}]$ over $K[\mathtt{rep}_{\widetilde{Q}}(\widetilde{\mathbf{d}})]$ and reducing modulo $\ker(\sigma^{\#})$ yields a $\mathbb{Z}^{d}$-graded free resolution of $K[\overline{\mathcal{O}}]$ over $K[\mathtt{rep}_Q(\mathbf{d})]$. The proof of this fact is the same as the equioriented case [MS05, Prop. 17.31], essentially relying on the fact that type A orbit closures are Cohen-Macaulay [BZ02] (see also [KR15, Prop. 6.1]). Since the K-polynomial can be computed as an alternating sum of K-polynomials of free modules in a resolution, this reduces the proposition to the case of free modules (i.e. $\overline{\mathcal{O}} = \mathtt{rep}_Q(\mathbf{d})$), which is exactly the statement that (5.13) is compatible with *sub*. ☐

5.2. **Ratio formula.** Now we can obtain formulas for K-polynomials of quiver loci in arbitrary orientation. First, we need to associate the same combinatorial objects to these orbit closures that were used in the bipartite case.

Definition 5.15. Let $\overline{\mathcal{O}} \subseteq \mathtt{rep}_Q(\mathbf{d})$ be an orbit closure. The *rank array* of $\overline{\mathcal{O}}$ is defined as the rank array $\mathbf{r}$ of $\overline{\pi^{-1}(\mathcal{O})}$. The *Zelevinsky permutation* of $\overline{\mathcal{O}}$ is the Zelevinsky permutation $v(\mathbf{r})$ of $\overline{\pi^{-1}(\mathcal{O})}$. ☐

As an immediate consequence of this definition and Proposition 5.14, we get a ratio formula for arbitrary orientation.

Theorem 5.16 (Ratio formula)**.** *Let $\overline{\mathcal{O}} \subseteq \mathtt{rep}_Q(\mathbf{d})$ be an orbit closure with rank array $\mathbf{r}$. Then we have*

$$\mathcal{K}_{\mathtt{rep}_Q(\mathbf{d})}(\overline{\mathcal{O}};\mathbf{s},\mathbf{t}) = sub\left(\frac{\mathfrak{G}_{v(\mathbf{r})}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})}{\mathfrak{G}_{v_*}(\mathbf{t},\mathbf{s};\mathbf{s},\mathbf{t})}\right), \tag{5.17}$$

and the same formula holds for multidegree upon replacing Grothendieck polynomials with Schubert polynomials.

5.3. **The pipe formula.** Define a partition into "good" and "bad" pipe dreams

$$\text{Pipes}(v_0, v(\mathbf{r})) = G \cup B$$

where G consists of pipe dreams P such that all +s of $P \setminus P_*$ are in blocks of the snake corresponding to original arrows of Q, and B consists of those with at least one + in an inverted block. In Figure 6, we see the row and column labels of a pipe dream from Figure 4 after applying *sub*.

Theorem 5.18 (Pipe formula)**.** *Let $\overline{\mathcal{O}} \subseteq \mathtt{rep}_Q(\mathbf{d})$ be an orbit closure with rank array $\mathbf{r}$. The K-polynomial $\mathcal{K}_{\mathtt{rep}_Q(\mathbf{d})}(\overline{\mathcal{O}};\mathbf{s},\mathbf{t})$ is given by applying sub to (3.8). Furthermore, the sum can be simplified to only summing over the subset $G \subseteq \text{Pipes}(v_0, v(\mathbf{r}))$. The analogous statement holds for the multidegree $\mathcal{C}_{\mathtt{rep}_Q(\mathbf{d})}(\overline{\mathcal{O}};\mathbf{s},\mathbf{t})$ and (3.9), and the sum can be taken over reduced pipe dreams in G.*

Proof. We only discuss the K-polynomial case, the multidegree case following from this as usual. As before, the first statement is immediate from Proposition 5.14. For the second statement, we need to show that the sum of terms indexed by B is zero. This can be seen by first fixing consideration of one inverted block of the snake region, say corresponding to a β-type arrow (α-type is similar), and partition B into subsets $\{B_i\}$ for which all elements in each B_i have the same configuration of +s outside of the designated block. Then further partition each B_i into subsets $B_i(w)$ indexed by permutations $w \neq 1$, such that for each $P \in B_i(w)$, the Demazure product of the restriction of P to the designated block is w. Then

$P_1 =$



FIGURE 6. Pipe dream from Figure 4 with inverted arrows

the sum of terms in the pipe formula over $B_i(w)$ has a factor $\mathfrak{G}_w(\mathbf{t}^i;\mathbf{s}^i)$ before applying *sub*, by [BR04, Theorem 2.1], which is sent to zero by *sub* by [BR04, Corollary 2.4] (see (5.21) below). Repeating over all inverted blocks yields the result. ☐

Remark 5.19. The last statement of the theorem gives a simplification of existing formulas in the literature for the equioriented case. Namely, the sums of [KMS06, "Pipe formula" p. 236] for multidegrees and [Mil05, Theorem 3] for K-polynomials can be taken over fewer pipe dreams (those with no +s on the block antidiagonal, in the setup of those papers). ☐

5.4. **The component formula.** In this section we give the component formula in arbitrary orientation. The formula for multidegrees was proven by Buch and Rimányi [BR07, Theorem 1]; the K-theoretic version was conjectured by them [BR07, Conjecture 1]. Note that minimal lacing diagrams for Q can be naturally identified with the subset of minimal lacing diagrams for $\widetilde{Q}$ which have no crossings at inverted arrows (equivalently, the identity permutation assigned to inverted arrows), and the same is true for K-theoretic lacing diagrams. So we can extend the definitions in (4.33) and (4.34) to get polynomials $\mathfrak{G}_{\mathbf{w}}(\mathbf{t};\mathbf{s})$ and $\mathfrak{S}_{\mathbf{w}}(\mathbf{t};\mathbf{s})$ associated to lacing diagrams of arbitrarily oriented Q.

Theorem 5.20 (Component formula). *Let $\overline{\mathcal{O}} \subseteq \mathtt{rep}_Q(\mathbf{d})$ be an orbit closure with rank array $\mathbf{r}$. Then the K-polynomial $\mathcal{K}_{\mathtt{rep}_Q(\mathbf{d})}(\overline{\mathcal{O}};\mathbf{s},\mathbf{t})$ is given by applying sub to* (4.38)*, and furthermore the sum can be restricted to K-theoretic lacing diagrams for $\overline{\mathcal{O}}$. The resulting sum is the same (up to differences in convention) as the formula in* [BR07, Conjecture 1]*. The analogous result holds for multidegrees, with summing over minimal lacing diagrams for $\overline{\mathcal{O}}$.*

Proof. As before we only explain the K-theoretic case. Applying Proposition 5.14 to (4.38), we *a priori* have sum over lacing diagrams for the $\widetilde{Q}$ quiver locus $\overline{\pi^{-1}(\mathcal{O})}$. Let δ be an inverted arrow of $\widetilde{Q}$. If $\mathbf{w}$ is a K-theoretic lacing diagram for $\overline{\pi^{-1}(\mathcal{O})}$, then the factor of $sub(\mathfrak{G}_{\mathbf{w}}(\mathbf{t};\mathbf{s}))$ corresponding to δ is of the form $\mathfrak{G}_w(\mathbf{t};\mathbf{t})$ for some permutation w. But we have, for example by [BR04, Cor. 2.4], that

$$\mathfrak{G}_w(\mathbf{t};\mathbf{t}) = \begin{cases} 1 & \text{if } w = id, \\ 0 & \text{if } w \neq id. \end{cases} \tag{5.21}$$

Applying this reasoning to all inverted arrows, we see that the only summands of (4.38) which survive after applying *sub* have no crossings over inverted arrows, thus are indexed by K-theoretic lacing diagrams associated to $\overline{\mathcal{O}}$. Furthermore the factors of each summand corresponding to inverted arrows of $\widetilde{Q}$ are just 1, so the summand indexed by such $\mathbf{w}$ agrees with the "intrinsic" definition of the summand associated to $\mathbf{w}$ in [BR07]. ☐



Figure 7. Minimal lacing diagrams for Q from contracting inverted arrows

Example 5.22. We take dimension vector $\mathbf{d} = (0, 2, 2, 3, 2, 1)$ for the quiver Q in (5.2), where vertices are read left to right, to continue our running example. Then, ignoring the vertex where $\mathbf{d}$ is 0, the associated bipartite quiver $\widetilde{Q}$ is the same as (2.4) in the running example from previous sections. The bipartite lacing diagrams of Figure 3 lie in the open subset $U \subset \mathtt{rep}_{\widetilde{Q}}(\widetilde{\mathbf{d}})$ of Theorem 5.7, and thus correspond to an orbit in $\mathtt{rep}_Q(\mathbf{d})$. In Figure 7, we see the corresponding Q-lacing diagrams and their extensions for this orbit, obtained by contracting inverted arrows in the lacing diagram from Figure 3 (i.e., applying π). ☐

References

[ADF85] S. Abeasis and A. Del Fra. Degenerations for the representations of a quiver of type A_m. *J. Algebra*, 93(2):376–412, 1985.

[AJS94] H. H. Andersen, J. C. Jantzen, and W. Soergel. Representations of quantum groups at a pth root of unity and of semisimple groups in characteristic p: independence of p. *Astérisque*, (220):321, 1994.

[All14] Justin Allman. Grothendieck classes of quiver cycles as iterated residues. *Michigan Math. J.*, 63(4):865–888, 2014.

[ASS06] Ibrahim Assem, Daniel Simson, and Andrzej Skowroński. *Elements of the representation theory of associative algebras. Vol. 1*, volume 65 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2006. Techniques of representation theory.

[BB93] Nantel Bergeron and Sara Billey. RC-graphs and Schubert polynomials. *Experiment. Math.*, 2(4):257–269, 1993.

[BF99] Anders Skovsted Buch and William Fulton. Chern class formulas for quiver varieties. *Invent. Math.*, 135(3):665–687, 1999.

[BFR05] Anders S. Buch, László M. Fehér, and Richárd Rimányi. Positivity of quiver coefficients through Thom polynomials. *Adv. Math.*, 197(1):306–320, 2005.

[Bil99] Sara C. Billey. Kostant polynomials and the cohomology ring for G/B. *Duke Math. J.*, 96(1):205–224, 1999.

[Bon98] Klaus Bongartz. Some geometric aspects of representation theory. In *Algebras and modules, I (Trondheim, 1996)*, volume 23 of *CMS Conf. Proc.*, pages 1–27. Amer. Math. Soc., Providence, RI, 1998.

[BR04] Anders S. Buch and Richárd Rimányi. Specializations of Grothendieck polynomials. *C. R. Math. Acad. Sci. Paris*, 339(1):1–4, 2004.

[BR07] Anders Skovsted Buch and Richárd Rimányi. A formula for non-equioriented quiver orbits of type *A*. *J. Algebraic Geom.*, 16(3):531–546, 2007.

[Buc02] Anders Skovsted Buch. Grothendieck classes of quiver varieties. *Duke Math. J.*, 115(1):75–103, 2002.

[Buc05] Anders Skovsted Buch. Alternating signs of quiver coefficients. *J. Amer. Math. Soc.*, 18(1):217–237 (electronic), 2005.

[Buc08] Anders Skovsted Buch. Quiver coefficients of Dynkin type. *Michigan Math. J.*, 57:93–120, 2008. Special volume in honor of Melvin Hochster.

[BZ02] Grzegorz Bobiński and Grzegorz Zwara. Schubert varieties and representations of Dynkin quivers. *Colloq. Math.*, 94(2):285–309, 2002.

[Eis95] David Eisenbud. *Commutative algebra, with a view toward algebraic geometry*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.

[FK94] Sergey Fomin and Anatol N. Kirillov. Grothendieck polynomials and the Yang-Baxter equation. In *Formal power series and algebraic combinatorics/Séries formelles et combinatoire algébrique*, pages 183–189. DIMACS, Piscataway, NJ, 1994.

[FK96] Sergey Fomin and Anatol N. Kirillov. The Yang-Baxter equation, symmetric functions, and Schubert polynomials. In *Proceedings of the 5th Conference on Formal Power Series and Algebraic Combinatorics (Florence, 1993)*, volume 153, pages 123–143, 1996.

[FL94] William Fulton and Alain Lascoux. A Pieri formula in the Grothendieck ring of a flag bundle. *Duke Math. J.*, 76(3):711–729, 1994.

[FR02] László Fehér and Richárd Rimányi. Classes of degeneracy loci for quivers: the Thom polynomial point of view. *Duke Math. J.*, 114(2):193–213, 2002.

[Ful92] William Fulton. Flags, Schubert polynomials, degeneracy loci, and determinantal formulas. *Duke Math. J.*, 65(3):381–420, 1992.

[Gra02] William Graham. Equivariant K-theory and Schubert varieties. Preprint, 2002.

[HZ14] B. Huisgen-Zimmermann. Fine and coarse moduli spaces in the representation theory of finite dimensional algebras. In *Expository lectures on representation theory*, volume 607 of *Contemp. Math.*, pages 1–34. Amer. Math. Soc., Providence, RI, 2014.

[KM04] Allen Knutson and Ezra Miller. Subword complexes in Coxeter groups. *Adv. Math.*, 184(1):161–176, 2004.

[KM05] Allen Knutson and Ezra Miller. Gröbner geometry of Schubert polynomials. *Ann. of Math. (2)*, 161(3):1245–1318, 2005.

[KMS06] Allen Knutson, Ezra Miller, and Mark Shimozono. Four positive formulae for type A quiver polynomials. *Invent. Math.*, 166(2):229–325, 2006.

[Knu] Allen Knutson. Frobenius splitting and Möbius inversion. `arxiv:0902.1930v1`.

[KR15] Ryan Kinser and Jenna Rajchgot. Type A quiver loci and Schubert varieties. *J. Commut. Algebra*, 7(2):265–301, 2015.

[KS11] Maxim Kontsevich and Yan Soibelman. Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson-Thomas invariants. *Commun. Number Theory Phys.*, 5(2):231–352, 2011.

[LM98] V. Lakshmibai and Peter Magyar. Degeneracy schemes, quiver schemes, and Schubert varieties. *Internat. Math. Res. Notices*, (12):627–640, 1998.

[LS82] Alain Lascoux and Marcel-Paul Schützenberger. Structure de Hopf de l'anneau de cohomologie et de l'anneau de Grothendieck d'une variété de drapeaux. *C. R. Acad. Sci. Paris Sér. I Math.*, 295(11):629–633, 1982.

[Lus90] G. Lusztig. Canonical bases arising from quantized enveloping algebras. *J. Amer. Math. Soc.*, 3(2):447–498, 1990.

[Mil05] Ezra Miller. Alternating formulas for K-theoretic quiver polynomials. *Duke Math. J.*, 128(1):1–17, 2005.

[MR85] V. B. Mehta and A. Ramanathan. Frobenius splitting and cohomology vanishing for Schubert varieties. *Ann. of Math. (2)*, 122(1):27–40, 1985.

[MS05] Ezra Miller and Bernd Sturmfels. *Combinatorial commutative algebra*, volume 227 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005.

[Rim] Richárd Rimányi. On the cohomological Hall algebra of Dynkin quivers. `arxiv:1303.3399`.

[Rim14] R. Rimányi. Quiver polynomials in iterated residue form. *J. Algebraic Combin.*, 40(2):527–542, 2014.

[Rin90] Claus Michael Ringel. Hall algebras and quantum groups. *Invent. Math.*, 101(3):583–591, 1990.

[RZ13] Christine Riedtmann and Grzegorz Zwara. Orbit closures and rank schemes. *Comment. Math. Helv.*, 88(1):55–84, 2013.

[Sch14] Ralf Schiffler. *Quiver representations*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer, Cham, 2014.

[Wil06] Matthieu Willems. K-théorie équivariante des tours de Bott. Application à la structure multiplicative de la K-théorie équivariante des variétés de drapeaux. *Duke Math. J.*, 132(2):271–309, 2006.

[WY08] Alexander Woo and Alexander Yong. Governing singularities of Schubert varieties. *J. Algebra*, 320(2):495–520, 2008.

[WY12] Alexander Woo and Alexander Yong. A Gröbner basis for Kazhdan-Lusztig ideals. *Amer. J. Math.*, 134(4):1089–1137, 2012.

[Yon05] Alexander Yong. On combinatorics of quiver component formulas. *J. Algebraic Combin.*, 21(3):351–371, 2005.

[Zel85] A. V. Zelevinsky. Two remarks on graded nilpotent classes. *Uspekhi Mat. Nauk*, 40(1(241)):199–200, 1985.

Department of Mathematics, University of Iowa, Iowa City, Iowa 52242, USA
E-mail address, Ryan Kinser: `ryan-kinser@uiowa.edu`

Cornell University, Department of Mathematics, Ithaca, NY, USA
E-mail address, Allen Knutson: `allenk@math.cornell.edu`

University of Michigan, Department of Mathematics, Ann Arbor, MI, USA
E-mail address, Jenna Rajchgot: `rajchgot@umich.edu`